

Crop Report

For the Period June 30 to July 6, 2026

Published by the Ministry of Agriculture
ISSN 0701 7085
Report number 10, July 9, 2026

Rainfall continued across many areas of the province this week, with some regions also experiencing minor to severe hail. Producers are beginning to see the impacts of the excessive moisture received over the past two weeks, including yellowing and drowned out crops. Crop development is highly variable across the province depending on weather conditions in the area, with many reporting slower development. Regular rainfall events have also delayed spraying and haying operations.

The Calder area recorded the highest rainfall with 112 millimetres (mm), followed by the Glaslyn area with 103 mm. The Radville and Churchbridge areas received 83 mm and 80 mm, respectively.

While many areas received rainfall this week, the surplus topsoil moisture conditions percentage decreased slightly. The surplus topsoil moisture ratings continue to be relatively high across the province overall.

Cropland topsoil moisture is:

- 30 per cent surplus;
- 69 per cent adequate; and
- 1 per cent short.

Hayland topsoil moisture is:

- 23 per cent surplus;
- 76 per cent adequate; and
- 1 per cent short.

Pasture topsoil moisture is:

- 16 per cent surplus;
- 82 per cent adequate; and
- 2 per cent short.

Due to differing environmental conditions around the province, stages of crop development are varying.

- 2 per cent of winter cereals are at the stem elongation stage, 8 per cent are at flag leaf, 56 per cent are heading and 34 per cent are at the dough stage.
- 1 per cent of spring cereals are pre-emergent, 3 per cent at seedling, 21 per cent tillering, 27 per cent at stem elongation, 34 per cent are at flag leaf and 14 per cent are at the heading stage.

One year ago

Rainfall and warm temperatures in some areas were allowing crops to progress nicely, while other areas continued to experience dry conditions which was stressing crops and leading to rapid development. Crop staging varied throughout the province and within regions as a result of irregular rainfall and made spray timing challenging.

Follow the 2026 Crop Report on X
[@SKAgriculture](#)

For further information, contact Rebecca Hort, BSc, AAg,
Crops Extension Specialist, Extension Services Branch,
Toll Free: 1-866-457-2377 or 306-946-6888, Email: cropreport@gov.sk.ca.
Also available on saskatchewan.ca/crop-report.



Agriculture and
Agri-Food Canada

SCIC
SASKATCHEWAN CROP
INSURANCE CORPORATION

Saskatchewan!

Crop Report

For the Period June 30 to July 6, 2026

Published by the Ministry of Agriculture

ISSN 0701 7085

Report number 10, July 9, 2026

- 24 per cent of flax crops are at the seedling stage, 62 per cent are at stem elongation, 13 per cent are flowering and 1 per cent at the boll stage.
- 8 per cent of canola and mustard crops are at the seedling stage, 27 per cent are at rosette, 37 per cent are bolting, 27 per cent are flowering and 1 per cent are podded.
- 2 per cent of pulse crops are at the seedling stage, 52 per cent are at the vegetative stage, 43 per cent are flowering and 3 per cent are podded.

Due to wet conditions, most livestock producers are still waiting to cut hay. Currently, four per cent of first cut hay has been cut and two per cent has been baled or silaged provincially. Hay quality is reported at 13 per cent excellent, 66 per cent good, eight per cent fair and 13 per cent poor. Like last week, most producers are still hoping for warmer and drier conditions to get out in the field to cut hay. Pasture conditions for the province are reported at 38 per cent excellent, 55 per cent good and seven per cent fair.

Several types of crop damage occurred throughout the province last week. Many producers are still reporting flooding as causing the most crop damage in the province and are beginning to see the effects. Severe thunderstorms last week brought hail to some areas in the province and caused damage to some crops. Additionally, producers have noted that gophers, cabbage seedpod weevil and wind have also caused minor crop damage.

In-crop spraying has been delayed in some areas of the province due to the wet conditions. Producers are trying to get out in the field to continue in-crop herbicide spraying and are evaluating if spraying fungicides and insecticides are warranted this week.

For further information, contact Rebecca Hort, BSc, AAg,
Crops Extension Specialist, Extension Services Branch,
Toll Free: 1-866-457-2377 or 306-946-6888, Email: cropreport@gov.sk.ca.
Also available on saskatchewan.ca/crop-report.




Agriculture and
Agri-Food Canada

SCIC
SASKATCHEWAN CROP
INSURANCE CORPORATION

Saskatchewan!

Crop Report Regions & Census Divisions



Southeastern Saskatchewan:

- Census Division 1 – Carnduff, Estevan, Lampman, Redvers and Stoughton areas
- Census Division 2 – Avonlea, Fillmore, Minton, Radville and Weyburn areas
- Census Division 5 – Broadview, Esterhazy, Melville and Moosomin areas
- Census Division 6 – Belle Plaine, Cupar, Lumsden, Indian Head, Regina and Rouleau areas

Weather varied in the southeast as some areas began to see warm and sunny weather, allowing them to finish spraying, while other areas are still delayed from wet and cold weather.

The Radville area received the most rainfall for this week with 83 millimetres (mm), followed closely by Churchbridge area with 80 mm. Rainfall varied variable throughout the region.

Surplus topsoil moisture decreased slightly compared to last week, with more of the region now reporting adequate levels.

Cropland topsoil moisture is:

- 31 per cent surplus; and
- 69 per cent adequate.

Hayland topsoil moisture is:

- 23 per cent surplus;
- 76 per cent adequate; and
- 1 per cent short.

Pasture topsoil moisture is:

- 23 per cent surplus;
- 75 per cent adequate; and
- 2 per cent short.

Crop staging varies throughout the region.

- 5 per cent of winter cereals are at the stem elongation stage, 3 per cent at flag leaf and 82 percent heading and 10 per cent at the dough stage.
- 3 per cent of spring cereals are at the tillering stage, 17 per cent tillering, 20 per cent at stem elongation, 34 per cent at flag leaf and 26 per cent heading.
- 18 per cent of flax crops are at the seedling stage, 76 per cent at stem elongation and 6 per cent flowering.
- 9 per cent of canola and mustard crops are at the seedling stage, 24 per cent at rosette, 33 per cent bolting and 34 per cent flowering.
- 5 per cent of pulse crops are at the seedling stage, 41 per cent vegetative and 54 per cent flowering.

Haying season has slowly begun in the southeast with four per cent cut. First cut has been rated as 10 per cent excellent quality, 73 per cent good and 17 per cent fair. Pasture conditions have been rated as 52 per cent excellent, 32 per cent good, 15 per cent fair and one per cent poor.

The primary sources of crop damage this week were flooding, cabbage seedpod weevil, hail and gophers, with impacts ranging from minor to severe. Wind caused minor to moderate damage in some areas. Minor crop damage from various insects and disease has been reported. Continued scouting for these pests will ensure management actions can be taken if warranted.

Southwestern Saskatchewan:

- Census Division 3 – Assiniboia, Gravelbourg, Mankota, Ponteix and Rockglen areas
- Census Division 4 – Cadillac, Consul, Eastend, Maple Creek and Val Marie areas
- Census Division 7 – Beechy, Central Butte, Craik, Herbert, Hodgeville and Moose Jaw areas
- Census Division 8 – Cabri, Elrose, Fox Valley, Leader, Swift Current and Tompkins areas

Variable precipitation around the region has some of the crops progressing well and others seeing some yellowing spots from excess moisture. Some producers are concerned about root rot in their lentil crop due to a few wet areas.

Smaller amounts of rainfall were received across the region this week compared to last week. The most rainfall was reported in the Lacadena area with 35 millimetres (mm) and the Mossbank area receiving 30 mm.

Topsoil moisture continues to be adequate throughout the southwest region, with small areas reporting surplus moisture.

Cropland topsoil moisture is:

- 14 per cent surplus;
- 85 per cent adequate; and
- 1 per cent short.

Hayland topsoil moisture is:

- 7 per cent surplus;
- 89 per cent adequate; and
- 4 per cent short.

Pasture topsoil moisture is:

- 10 per cent surplus;
- 85 per cent adequate; and
- 5 per cent short.

Crop staging is variable throughout the region with some crops still at the seedling stage and others beginning to develop pods.

- 1 per cent of winter cereals are at the flag leaf stage, 52 per cent heading and 47 per cent at the dough stage.
- 2 per cent of spring cereals are pre-emergent, 9 per cent tillering, 24 per cent at stem elongation, 41 per cent at flag leaf and 24 per cent heading.

- 15 per cent of flax crops are at the seedling stage, 51 per cent at stem elongation, 30 per cent flowering and 4 per cent at the boll stage.
- 12 per cent of canola and mustard crops are at the rosette stage, 31 per cent bolting, 52 per cent flowering and 5 per cent podded.
- 1 per cent of pulse crops are at the seedling stage, 44 per cent vegetative, 50 per cent flowering and 5 per cent podded.

Hay has been reported as eight per cent cut and four per cent baled or silage. Quality has been rated as 24 per cent excellent, 71 per cent good and five per cent fair. Pasture conditions have been rated as 33 per cent excellent, 58 per cent good and nine per cent fair.

The main sources of crop damage from this week included minor to severe reports of hail, cabbage seedpod weevil, gophers and root rots in pulses. Flooding, alfalfa weevil, aphids and sclerotinia caused some minor damage in the southwest. Several insect pests caused some minor damage as well.

East-Central Saskatchewan:

- Census Division 9 – Calder, Canora, Pelly, Preeceville, Sheho and Yorkton areas
- Census Division 10 – Foam Lake, Kelliher, Leroy, Raymore and Wadena areas
- Census Division 11 – Davidson, Colonsay, Langham, Lanigan, Nokomis, Outlook and Saskatoon areas

There was patchy precipitation throughout the east-central region this past week. Some producers got a well needed break from rain; however, flooding is still a concern with many fields extremely wet and holding water in low spots. Producers are trying to get spraying done but still have some fields too wet to spray.

Many areas of the east-central region received rainfall this week. The most rainfall was 112 millimetres (mm) and 80 mm that fell in the Calder area. There was notable rainfall recorded in other areas, with several other parts of the region receiving up to 65 mm.

There is still a significant amount of surplus topsoil moisture this week in this region.

Cropland topsoil moisture is:

- 45 per cent surplus;
- 54 per cent adequate; and
- 1 per cent short.

Hayland topsoil moisture is:

- 43 per cent surplus;
- 55 per cent adequate; and
- 2 per cent short.

Pasture topsoil moisture is:

- 40 per cent surplus;
- 58 per cent adequate; and
- 2 per cent short.

Crop staging is variable throughout the east-central region with a few acres still having pre-emergent and seedling stage crops and some crops beginning to flower.

- 5 per cent of winter cereals are at the stem elongation stage, 8 per cent are at flag leaf, 84 per cent are heading, and 3 per cent are at the dough stage.
- 1 per cent of spring cereals are pre-emergent, 9 per cent are at the seedling stage, 34 per cent are tillering, 24 per cent are at stem elongation, 22 per cent are at flag leaf and 10 per cent are heading.
- 30 per cent of flax crops are at the seedling stage, 49 per cent are at stem elongation and 21 per cent are flowering.
- 1 per cent of canola and mustard are pre-emergent, 15 per cent are at the seedling stage, 36 per cent at rosette, 31 per cent are bolting and 17 per cent are flowering.
- 1 per cent of pulse crops are at the seedling stage, 54 per cent at vegetative stage and 45 per cent are flowering.

Haying season has just begun in the east-central region with three per cent cut. Quality has been rated as 53 per cent excellent, 30 per cent good, 16 per cent fair and one per cent poor. Pasture conditions have been rated as 44 per cent excellent, 50 per cent good, four per cent fair and two per cent poor.

The main cause of crop damage in the east-central region this week was flooding, ranging from minor to severe, with producers seeing the effects of the consistent moisture the past few weeks. Hail, wind, gophers and cabbage seedpod weevil caused minor to moderate damage in some areas. Flea beetles and root rots are showing some moderate damage in areas. Producers and agronomists are keeping an eye on the minor damage being caused by disease in the east-central region.

West-Central Saskatchewan:

- Census Division 12 – Biggar, Delisle, Rosetown and Sonningdale areas
- Census Division 13 – Cut Knife, Kerrobert, Kindersley, Macklin, Plenty and Wilkie areas

Spotty precipitation occurred in the west-central region with some areas receiving trace amounts of rain with a few receiving over an inch. Moisture varies in the region with some areas still being too wet but most areas have adequate amounts of moisture. There have been localized hail events with minimal crop damage reported. Producers have started fungicide applications on pulses and canola.

The most rain fell in the Vanscoy region with 75 millimetres (mm) followed by the Buffalo region with 45 mm. Some areas of the west-central region received trace amounts of rain while others received over an inch.

Topsoil moisture continues to be adequate throughout the west-central region, with small areas reporting surplus moisture.

Cropland topsoil moisture is:

- 12 per cent surplus; and
- 88 per cent adequate.

Hayland topsoil moisture is:

- 9 per cent surplus; and
- 91 per cent adequate.

Pasture topsoil moisture is:

- 4 per cent surplus; and
- 96 per cent adequate.

Various crop stages are reported throughout the region with some at seedling and others beginning to develop pods.

- 2 per cent of winter cereals are at the flag leaf stage, 76 per cent are heading and 22 per cent are at the dough stage.
- 2 per cent of spring cereals are at the seedling stage, 17 per cent are tillering, 30 per cent at stem elongation, 41 per cent at flag leaf and 10 per cent are heading.
- 6 per cent of flax crops are at the seedling stage, 84 per cent are at stem elongation and 10 per cent are flowering.
- 21 per cent of canola and mustard crops are at the rosette stage, 46 per cent are bolting and 33 per cent are flowering.
- 1 per cent of pulse crops are at the seedling stage, 61 per cent are at the vegetative stage, 37 per cent are flowering and 1 per cent are podded.

Haying season has just begun in the west-central region with three per cent cut. Quality has been rated as 23 per cent excellent, 76 per cent good and one per cent fair. Pasture conditions have been rated as 40 per cent excellent, 55 per cent good and five per cent fair.

Minor to moderate crop damage from flooding and hail damage was reported in localized areas in the region. Wind and gopher damage was reported as minor. Limited disease damage was reported in some of the region with producers and agronomists continuing to scout for these pests.

Northeastern Saskatchewan:

- Census Division 14 – Choiceland, Hudson Bay, Kelvington, Melfort, and Nipawin areas
- Census Division 15 – Cudworth, Humboldt, Kinistino, Prince Albert, Rosthern and St. Brieux areas

Conditions are variable in the northeast region as some areas continued to see excess rain this week, causing wet conditions, while other areas of the region have reported good growing conditions and crops advancing normally. Crops have begun to yellow in overly wet areas. Some producers have begun spraying fungicides.

Significant rainfall was seen in the Middle Lake and Archerwill areas with 76 millimetres (mm) and 71 mm respectively. The Rosthern area received 58 mm and Arborfield and Bruno areas both received 53 mm. There was notable rainfall recorded in other areas, with most of the region receiving significant amounts.

The surplus topsoil moisture is still relatively high in the northeast region.

Cropland topsoil moisture is:

- 44 per cent surplus;
- 55 per cent adequate; and
- 1 per cent short.

Hayland topsoil moisture is:

- 39 per cent surplus; and
- 61 per cent adequate.

Pasture topsoil moisture is:

- 36 per cent surplus; and
- 64 per cent adequate.

Variable crop staging is seen throughout the northeast with some canola and pulses beginning to flower.

- 41 per cent of winter cereals are at the flag leaf stage and 59 per cent are heading.
- 31 per cent of spring cereals are tillering, 41 per cent are at stem elongation, 25 per cent are at flag leaf and 3 per cent are heading.
- 12 per cent of flax crops are at the seedling stage, 86 per cent at stem elongation and 2 per cent flowering.
- 8 per cent of canola and mustard crops are at the seedling stage, 35 per cent are at the rosette stage, 43 per cent are bolting and 14 per cent are flowering.
- 3 per cent of pulse crops are at the seedling stage, 75 per cent are at the vegetative stage and 22 per cent are flowering.

Haying season has just begun with three per cent cut and three per cent baled or silaged. Quality has been rated as 33 per cent excellent and 67 per cent good. Pasture conditions have been rated as 50 per cent excellent, 47 per cent good and three per cent fair.

The main cause of crop damage in the northeast is flooding, ranging from minor to severe, depending on the area. Cabbage seedpod weevil has caused some minor and severe damage in areas of the region. Hail caused minor to moderate damage in some areas. Wind, gophers and flea beetles caused minor damage depending on the area.

Northwestern Saskatchewan:

- Census Division 16 – Blaine Lake, Canwood, North Battleford, Radisson and Spiritwood areas
- Census Division 17 – Glaslyn, Maidstone, Meadow Lake, Pierceland and St. Walburg areas

The northwest region experienced variable weather this week, including some isolated hail and heavy rainstorms. Crops in areas that were not affected by those storms are seeing good crop development and fungicides are being applied.

With the heavy rainstorms in the region, most areas of the northwest received significant amounts of rainfall this week. The most rainfall was recorded in the Glaslyn area with 103 millimetres (mm). Blaine Lake and other areas of the Parkdale area received 65 mm of rain. There was notable rainfall recorded in other areas, with most of the region receiving significant amounts.

Topsoil moisture continues to be adequate throughout the northwest region, with small areas reporting surplus moisture.

Cropland topsoil moisture is:

- 23 per cent surplus; and
- 77 per cent adequate.

Hayland topsoil moisture is:

- 15 per cent surplus; and
- 85 per cent adequate.

Pasture topsoil moisture is:

- 13 per cent surplus;
- 86 per cent adequate; and
- 1 per cent short.

Crop staging is variable throughout the region with pulses, winter cereals and flax the furthest along development.

- 100 per cent of winter cereal crops have been reported at the dough stage.
- 1 per cent of spring cereals are at the seedling stage, 16 per cent are tillering, 29 per cent are at stem elongation, 38 per cent are at flag leaf and 16 per cent are heading.
- 26 per cent of flax crops are at stem elongation and 74 per cent are flowering.
- 1 per cent of canola and mustard crops are at the seedling stage, 22 per cent are at rosette, 47 per cent are bolting and 30 per cent are flowering.
- 61 per cent of pulse crops are at the vegetative stage and 39 per cent are flowering.

Haying season has just begun with two per cent cut. Quality has been rated as 43 per cent excellent, 56 per cent good and one per cent fair. Pasture conditions have been rated as 46 per cent excellent, 52 per cent good and two per cent fair.

The main causes of crop damage was flooding, hail and gophers with damage ranging from minor to severe based on the area. Wind and cabbage seedpod weevil was reported to have caused minor to moderate damage. There has been limited minor damage caused by disease and insects in the northeast with agronomists and producers continuing scouting for pests.

Crop Staging Tables-June 30th to July 6th, 2026

Winter Cereals	Tillering	Stem Elongation	Flag Leaf	Heading	Dough	Ripe
South East	0%	5%	3%	82%	10%	0%
South West	0%	0%	1%	52%	47%	0%
East Central	0%	5%	8%	84%	3%	0%
West Central	0%	0%	2%	76%	22%	0%
North East	0%	0%	41%	59%	0%	0%
North West	0%	0%	0%	0%	100%	0%
Provincial	0%	2%	8%	56%	34%	0%

Spring Cereals	Pre Emerging	Seedling	Tillering	Stem Elongation	Flag Leaf	Heading	Dough	Ripe
South East	0%	3%	17%	20%	34%	26%	0%	0%
South West	2%	0%	9%	24%	41%	24%	0%	0%
East Central	1%	9%	34%	24%	22%	10%	0%	0%
West Central	0%	2%	17%	30%	41%	10%	0%	0%
North East	0%	0%	31%	41%	25%	3%	0%	0%
North West	0%	1%	16%	29%	38%	16%	0%	0%
Provincial	1%	3%	21%	27%	34%	14%	0%	0%

Flax	Pre Emerging	Seedling	Stem Elongation	Flowering	Boll	Ripe
South East	0%	18%	76%	6%	0%	0%
South West	0%	15%	51%	30%	4%	0%
East Central	0%	30%	49%	21%	0%	0%
West Central	0%	6%	84%	10%	0%	0%
North East	0%	12%	86%	2%	0%	0%
North West	0%	0%	26%	74%	0%	0%
Provincial	0%	24%	62%	13%	1%	0%

Canola and Mustard	Pre Emerging	Seedling	Rosette	Bolting	Flowering	Podded	Ripe
South East	0%	9%	24%	33%	34%	0%	0%
South West	0%	0%	12%	31%	52%	5%	0%
East Central	1%	15%	36%	31%	17%	0%	0%
West Central	0%	0%	21%	46%	33%	0%	0%
North East	0%	8%	35%	43%	14%	0%	0%
North West	0%	1%	22%	47%	30%	0%	0%
Provincial	0%	8%	27%	37%	27%	1%	0%

Pulse Crops	Pre Emerging	Seedling	Vegetative	Flowering	Podded	Ripe
South East	0%	5%	41%	54%	0%	0%
South West	0%	1%	44%	50%	5%	0%
East Central	0%	1%	54%	45%	0%	0%
West Central	0%	1%	61%	37%	1%	0%
North East	0%	3%	75%	22%	0%	0%
North West	0%	0%	61%	39%	0%	0%
Provincial	0%	2%	52%	43%	3%	0%

Weekly Rainfall Summary

(reported in millimeters)

1 inch=25 mm

for the period from June 30 to July 6, 2026

Census Division	RM No.	RM Name	Past Week	Since 1-Apr	Census Division	RM No.	RM Name	Past Week	Since 1-Apr	Census Division	RM No.	RM Name	Past Week	Since 1-Apr
1	3	ENNISKILLEN	52	281	8	138	WEBB	18	224	14	366	KELVINGTON	24	212
1	4	COALFIELDS	40	252	8	139	GULL LAKE	22	202	14	367	PONASS LAKE	52	271
1	32	RECIPROCITY	43	249	8	168	RIVERSIDE	20	193	14	394	HUDSON BAY	27	204
1	34	BROWNING	N/A	162	8	228	LACADENA	35	232	14	395	PORCUPINE	32	233
1	94	HAZELWOOD	N/A	0	8	229	MIRY CREEK	N/A	143	14	397	BARRIER VALLEY	N/A	166
1	95	GOLDEN WEST	37	225	8	231	HAPPYLAND	12	157	14	428	STAR CITY	39	176
1	2A	MOUNT PLEASANT	29	188	8	259	SNIPER LAKE	0	174	14	456	ARBORFIELD	53	201
2	10	HAPPY VALLEY	67	157	8	138A	WEBB	12	149	14	457	CONNAUGHT	30	67
2	38	LAURIER	83	232	8	228A	LACADENA	N/A	64	14	486	MOOSE RANGE	N/A	113
2	66	GRIFFIN	56	244	8	257A	MONET	5	188	14	487	NIPAWIN	33	163
2	67	WEYBURN	N/A	101	8	259A	SNIPER LAKE	N/A	141	14	488	TORCH RIVER	28	175
2	70	KEY WEST	N/A	12	9	241	CALDER	112	363	14	394A	HUDSON BAY	N/A	200
2	100	ELMSTHORPE	N/A	174	9	243	WALLACE	N/A	118	14	397A	BARRIER VALLEY	71	200
2	100A	ELMSTHORPE	38	213	9	245	GARRY	13	124	14	488A	TORCH RIVER	N/A	55
2	38A	LAURIER	N/A	96	9	273	SLIDING HILLS	N/A	275	14	488B	TORCH RIVER	N/A	5
3	73	STONEHENGE	5	179	9	274	GOOD LAKE	46	213	15	369	ST. PETER	46	193
3	74	WOOD RIVER	15	177	9	301	ST. PHILIPS	N/A	115	15	371	BAYNE	53	210
3	75	PINTO CREEK	4	121	9	305	INVERMAY	65	260	15	372	GRANT	27	161
3	76	AUVERGNE	9	162	9	331	LIVINGSTON	N/A	206	15	373	ABERDEEN	23	135
3	101	TERRELL	N/A	0	9	241A	CALDER	80	436	15	399	LAKE LENORE	45	232
3	102	LAKE JOHNSTON	30	173	9	245A	GARRY	59	186	15	400	THREE LAKES	76	278
3	106	WHISKA CREEK	11	178	10	246	ITUNA BON ACCORD	8	107	15	402	FISH CREEK	N/A	116
3	74A	KEY WEST	12	152	10	247	KELLROSS	37	172	15	429	FLETT'S SPRINGS	42	188
4	51	RENO	0	112	10	248	TOUCHWOOD	32	185	15	430	INVERGORDON	N/A	100
4	79	ARLINGTON	4	201	10	277	EMERALD	40	187	15	459	KINISTINO	25	199
4	110	PIAPOT	4	210	10	279	MOUNT HOPE	N/A	151	15	461	PRINCE ALBERT	40	159
4	77A	WISE CREEK	N/A	152	10	307	ELFROS	31	330	15	463	DUCK LAKE	46	176
4	78A	GRASSY CREEK	N/A	107	10	309	PRAIRIE ROSE	40	225	15	491	BUCKLAND	36	90
4	79A	ARLINGTON	3	226	10	336	SASMAN	63	282	15	520	PADDOCKWOOD	20	20
5	122	MARTIN	17	257	10	339	LEROY	28	200	15	521	LAKELAND	20	20
5	124	KINGSLEY	12	215	10	246A	ITUNA BON ACCORD	N/A	72	15	371A	BAYNE	20	154
5	151	ROCANVILLE	3	208	10	248A	TOUCHWOOD	36	312	15	371B	BAYNE	39	216
5	155	WOLSELEY	22	194	10	276A	FOAM LAKE	64	341	15	399A	LAKE LENORE	N/A	16
5	181	LANGENBURG	16	403	10	276B	FOAM LAKE	65	408	15	403A	ROSTHERN	26	181
5	183	FERTILE BELT	15	265	10	276C	FOAM LAKE	56	263	15	403C	ROSTHERN	58	211
5	211	CHURCHBRIDGE	54	286	10	276D	FOAM LAKE	28	155	15	461A	PRINCE ALBERT	N/A	99
5	213	SALTCOATS	48	316	10	277A	EMERALD	N/A	184	15	461B	PRINCE ALBERT	N/A	2
5	215	STANLEY	10	218	10	279A	MOUNT HOPE	N/A	156	16	434	BLAINE LAKE	65	77
5	125A	CHESTERFIELD	35	180	11	251	BIG ARM	0	134	16	435	REDBERRY	11	248
5	154A	ELCAPO	14	206	11	280	WRETFORD	N/A	83	16	436	DOUGLAS	27	119
5	183A	FERTILE BELT	N/A	275	11	283	ROSEDALE	46	244	16	437	NORTH BATTLEFORD	58	267
5	211A	CHURCHBRIDGE	80	303	11	284	RUDY	0	65	16	466	MEETING LAKE	N/A	131
6	127	FRANCIS	18	215	11	314	DUNDURN	27	188	16	467	ROUND HILL	N/A	95
6	128	LAJORD	31	111	11	344	CORMAN PARK	20	136	16	493	SHELLBROOK	31	117
6	130	REDBURN	64	233	11	282A	McCRANEY	60	150	16	494	CANWOOD	31	104
6	156	INDIAN HEAD	19	224	12	286	MILDEN	N/A	174	16	497	MEDSTEAD	57	191
6	160	PENSE	N/A	145	12	287	ST. ANDREWS	1	160	16	437A	NORTH BATTLEFORD	N/A	195
6	186	ABERNETHY	10	231	12	288	PLEASANT VALLEY	13	155	16	467A	ROUND HILL	39	178
6	190	DUFFERIN	21	175	12	316	HARRIS	10	128	17	468	MEOTA	60	143
6	216	TULLYMET	N/A	10	12	317	MARRIOTT	15	190	17	470	PAYNTON	1	44
6	217	LIPTON	64	284	12	345	VANSCOY	75	130	17	471	ELDON	25	215
6	219	LONGLAKETON	35	226	12	346	PERDUE	0	158	17	498	PARKDALE	103	237
6	221	SARNIA	11	199	12	347	BIGGAR	4	204	17	499	MERVIN	60	176
6	127B	FRANCIS	N/A	116	12	376	EAGLE CREEK	9	207	17	502	BRITANNIA	24	158
6	159A	SHERWOOD	14	131	12	377	GLENSIDE	7	190	17	561	LOON LAKE	26	126
6	159B	SHERWOOD	10	184	12	285A	FERTILE VALLEY	10	143	17	588	MEADOW LAKE	51	210
6	160A	PENSE	18	138	13	290	KINDERSLEY	N/A	127	17	622	BEAVER RIVER	28	190
6	190A	DUFFERIN	31	116	13	292	MILTON	32	318	17	498A	PARKDALE	65	199
6	190B	DUFFERIN	35	202	13	321	PRAIRIEDALE	18	228	17	501A	FRENCHMAN BUTTE	37	170
6	190C	DUFFERIN	48	246	13	350	MARIPOSA	30	195	17	501C	FRENCHMAN BUTTE	N/A	50
6	216A	TULLYMET	N/A	75	13	351	PROGRESS	10	214	17	561A	LOON LAKE	43	139
6	217A	LIPTON	N/A	2	13	379	REFORD	N/A	93					
6	219A	LONGLAKETON	34	223	13	382	EYE HILL	35	187					
6	219B	LONGLAKETON	N/A	60	13	409	BUFFALO	16	247					
6	220A	McKILLOP	9	161	13	440	HILLSDALE	N/A	173					
7	132	HILLSBOROUGH	21	288	13	442	MANITOU LAKE	31	205					
7	136	COULEE	10	151	13	292A	MILTON	11	171					
7	161	MOOSE JAW	18	137	13	320A	OAKDALE	10	124					
7	162	CARON	14	207	13	321A	PRAIRIEDALE	N/A	35					
7	165	MORSE	18	238	13	409A	BUFFALO	45	204					
7	193	EYEBROW	25	229										
7	223	HURON	0	183										
7	132A	HILLSBOROUGH	12	113										
7	136A	COULEE	N/A	0										
7	161A	MOOSE JAW	N/A	38										
7	162A	CARON	13	183										
7	222A	CRAIK	0	215										
7	223A	HURON	15	167										

Municipality No: A, B, C and D - more than one reporter



These precipitation amounts represent point locations within each municipality and do not necessarily reflect the whole R. M.

N/A indicates that rainfall was not reported for the week

Weekly Rainfall

from June 30 to July 6, 2026

Rainfall (mm)





NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.



Cumulative Rainfall

from April 1 to July 6, 2026

Rainfall (mm)




NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.



Pasture Conditions

from June 30 to July 6, 2026

Pasture Conditions



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.



Topsoil Moisture Conditions-for the period of June 30th to July 6th, 2026


Cropland				
Region	Surplus	Adequate	Short	Very Short
South East	31%	69%	0%	0%
South West	14%	85%	1%	0%
East Central	45%	54%	1%	0%
West Central	12%	88%	0%	0%
North East	44%	55%	1%	0%
North West	23%	77%	0%	0%
Provincial Average	30%	69%	1%	0%

Hayland				
Region	Surplus	Adequate	Short	Very Short
South East	23%	76%	1%	0%
South West	7%	89%	4%	0%
East Central	43%	55%	2%	0%
West Central	9%	91%	0%	0%
North East	39%	61%	0%	0%
North West	15%	85%	0%	0%
Provincial Average	23%	76%	1%	0%

Pasture				
Region	Surplus	Adequate	Short	Very Short
South East	23%	75%	2%	0%
South West	10%	85%	5%	0%
East Central	40%	58%	2%	0%
West Central	4%	96%	0%	0%
North East	36%	64%	0%	0%
North West	13%	86%	1%	0%
Provincial Average	16%	82%	2%	0%

Cropland Topsoil Moisture Conditions

from June 30 to July 6, 2026




NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.



Hay Topsoil Moisture Conditions

from June 30 to July 6, 2026




NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.



Pasture Topsoil Moisture Conditions

from June 30 to July 6, 2026




NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.




Maximum Temperature

from June 30 to July 6, 2026



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.




Data Sources:
 Temperature data - Saskatchewan Ministry of Environment (Wildfire Management Branch) and Environment Canada.
 Temperature data compiled and quality controlled by Agriculture and Agri-Food Canada
 IDW interpolation (power 3.5, fixed radius 300 km)
 Geomatics Services, Ministry of Agriculture


July 8, 2026

Minimum Temperature

from June 30 to July 6, 2026



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.



Projection: UTM Zone 13 Datum: NAD83



Data Sources:
 Temperature data - Saskatchewan Ministry of Environment (Wildfire Management Branch) and Environment Canada.
 Temperature data compiled and quality controlled by Agriculture and Agri-Food Canada
 IDW interpolation (power 3.5, fixed radius 300 km)
 Geomatics Services, Ministry of Agriculture

July 8, 2026