

Crop Report

For the Period July 28 to August 3, 2026

Published by the Ministry of Agriculture

ISSN 0701 7085

Report number 14, August 6, 2026

Provincial Crop Development			
Crop	% Ahead	% Normal	% Behind
Fall Cereals	4%	86%	10%
Spring Cereals	7%	71%	22%
Oilseeds	6%	65%	29%
Pulse Crops	10%	77%	13%
Perennial Forage	2%	83%	15%
Annual Forage	1%	78%	21%

Crops rapidly developed due to warm temperatures and dry conditions in many areas across the province this week. Early-seeded crops are nearing maturity in many areas, and rain likely would not benefit these crops. However, producers in other areas are hoping for rain soon to help with grain fill in later-seeded crops.

While scattered showers brought limited moisture in some regions, most received no precipitation. The Milden area received the most rainfall with 31 millimetres mm , followed by the Beaver River area with 28 mm and the nvermay area with 20 mm.

Despite the lack of rainfall, topsoil moisture is rated as adequate for over half of the province.

Cropland topsoil moisture is:

- 2 per cent surplus;
- 56 per cent adequate;
- 33 per cent short; and
- 9 per cent very short.

Hayland topsoil moisture is:

- 2 per cent surplus;
- 56 per cent adequate;
- 34 per cent short; and
- 8 per cent very short.

One year ago

Crops continued to develop rapidly due to hot and dry conditions across the province. Harvest had started in some areas as one per cent of crops had been taken off, mainly winter cereals and pulse crops. Although most of the pastures were in good to fair conditions, a notable portion were also in poor conditions due to limited precipitation in certain areas. Most producers were preparing for harvest by servicing equipment, hauling grain, and cleaning bins.

Follow the 2026 Crop Report on X
@SKAgriculture

<u>Harvest Progress in SK</u> Per Cent Combined Historical All Crops	
Aug 3/26	2
Aug 4/25	1
Aug 5/24	2
Aug 7/23	4
Aug 8/22	1
Aug 2/21	3
5 year avg. (2020-2024)	2
10 year avg. (2015-2024)	2

For further information, contact Augustine Osei, PhD, PAg,
Crops Extension Specialist, Extension Services Branch,
Toll Free: 1-866-457-2377 or 306-520-5761, Email: cropreport@gov.sk.ca.
Also available on the Ministry of Agriculture website at saskatchewan.ca/crop-report.



Agriculture and
Agri-Food Canada

SCIC
SASKATCHEWAN CROP
INSURANCE CORPORATION

Saskatchewan

Crop Report

For the Period July 28 to August 3, 2026

Published by the Ministry of Agriculture
ISSN 0701 7085
Report number 14, August 6, 2026

Pasture topsoil moisture is:

- 2 per cent surplus;
- 45 per cent adequate;
- 41 per cent short; and
- 12 per cent very short

The hot and dry weather conditions have accelerated crop development, with most crops being rated as normal for this time of year.

- 4 per cent of fall cereals are reported as ahead, 86 per cent as normal and 10 per cent behind.
- 7 per cent of spring cereals are reported as ahead, 71 per cent as normal and 22 per cent behind.
- 6 per cent of oilseeds are reported as ahead, 65 per cent as normal and 29 per cent behind.
- 10 per cent of pulse crops are reported as ahead, 77 per cent as normal and 13 per cent behind.
- 2 per cent of perennial forages are reported as ahead, 83 per cent as normal and 15 per cent behind.
- 1 per cent of annual forages are reported as ahead, 78 per cent as normal and 21 per cent behind.

The dry conditions for much of the season in certain areas have led to some producers starting to harvest early-seeded crops and winter cereals. Provincially, harvest is two per cent complete with most of this progress occurring in the southwest.

Winter cereals harvested:

- Winter wheat: 7 per cent; and
- Fall rye: 21 per cent.

Spring cereals harvested:

- Triticale: 37 per cent;
- Barley: 1 per cent;
- Spring wheat: 1 per cent; and
- Oats: 2 per cent.

Many pastures are in good to fair condition, but a small portion is rated as in poor condition due to limited precipitation in certain areas. Currently, six per cent of pastures are in excellent condition, 53 per cent are good, 33 per cent are fair, seven per cent are poor, and one per cent is in very poor condition.

Crop damage was reported in areas experiencing hot temperatures paired with dry conditions, with some areas reporting minor to moderate damage. There were also reports of wind, grasshoppers and bertha armyworms causing minor crop damage in certain areas.

For further information, contact Augustine Osei, PhD, PAg,
Crops Extension Specialist, Extension Services Branch,
Toll Free: 1-866-457-2377 or 306-520-5761, Email: cropreport@gov.sk.ca.
Also available on the Ministry of Agriculture website at saskatchewan.ca/crop-report.



**Agriculture and
Agri-Food Canada**

SCIC
SASKATCHEWAN CROP
INSURANCE CORPORATION

Saskatchewan

Crop Report

For the Period July 28 to August 3, 2026

Published by the Ministry of Agriculture
ISSN 0701 7085
Report number 14, August 6, 2026

As producers prepare for and begin harvest, there will be increased farm equipment traffic on public roadways. Drivers are reminded to have extra patience and caution when encountering large equipment on the road. Additionally, farm equipment operating in dry fields can present a fire risk. Producers are encouraged to take precautions to mitigate fire risk and ensure harvest efforts are conducted safely.

For crop or livestock questions, producers are encouraged to call the Agriculture Knowledge Centre toll free at 1-866-457-2377.

This can be a stressful time of year for producers as weather conditions can be unpredictable. The Farm Stress Line can help by providing support for producers toll free at 1-800-667-4442.

A complete, printable version of the Crop Report is available online .

Follow the 2026 Crop Report on X at @SKAgriculture.

For more information, contact:

Augustine Osei
Saskatchewan Agriculture
Weyburn
Phone: 306-520-5761
Email: augustine.osei@gov.sk.ca

For further information, contact Augustine Osei, PhD, PAg,
Crops Extension Specialist, Extension Services Branch,
Toll Free: 1-866-457-2377 or 306-520-5761, Email: cropreport@gov.sk.ca.
Also available on the Ministry of Agriculture website at saskatchewan.ca/crop-report.



Agriculture and
Agri-Food Canada



SCIC
SASKATCHEWAN CROP
INSURANCE CORPORATION

Saskatchewan

Harvest Progress


from April 1 to August 3, 2026

Harvest Progress



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Crop Report Regions & Census Divisions



Southeastern Saskatchewan:

- Census Division 1 – Carnduff, Estevan, Lampman, Redvers and Stoughton areas
- Census Division 2 – Avonlea, Fillmore, Minton, Radville and Weyburn areas
- Census Division 5 – Broadview, Esterhazy, Melville and Moosomin areas
- Census Division 6 – Belle Plaine, Cupar, Lumsden, Indian Head, Regina and Rouleau areas

Hot, dry weather continues to advance crop development in the southeast. Producers would welcome rain for later seeded crops and pastures.

Like much of the province, there were only a few scattered showers in a few areas in the region, providing very little precipitation. The highest precipitation received in the region was only 9 millimeters (mm) in the Indian Head area with the Reciprocity and Kingsley areas receiving 7 mm. This was closely followed by the Mount Pleasant and Langenburg areas receiving 5 mm of precipitation. A large part of the region did not receive any precipitation at all, impacting topsoil moisture in the region.

Cropland topsoil moisture is:

- 1 per cent surplus;
- 60 per cent adequate;
- 38 per cent short; and
- 1 per cent very short.

Hayland topsoil moisture is:

- 1 per cent surplus;
- 53 per cent adequate;
- 44 per cent short; and
- 2 per cent very short.

Pasture topsoil moisture is:

- 2 per cent surplus;
- 46 per cent adequate;
- 47 per cent short; and
- 5 per cent very short.

Southeast Saskatchewan Crop Development			
Crop	% Ahead	% Normal	% Behind
Fall Cereals	6%	89%	5%
Spring Cereals	10%	76%	14%
Oilseeds	8%	74%	18%
Pulse Crops	10%	78%	12%
Perennial Forage	4%	90%	6%
Annual Forage	5%	89%	6%

Warm conditions have caused crop development to accelerate, offsetting earlier delays, with most crops reported to be at normal stage.

Crop development for this time of year in the region is rated as follows:

- 6 per cent of fall cereals are reported as ahead, 89 per cent as normal and 5 per cent behind.
- 10 per cent of spring cereals are reported as ahead, 76 per cent as normal and 14 per cent behind.
- 8 per cent of oilseeds are reported as ahead, 74 per cent as normal and 18 per cent behind.
- 10 per cent of pulse crops are reported as ahead, 78 per cent as normal and 12 per cent behind.
- 4 per cent of perennial forage are reported as ahead, 90 per cent as normal and 6 per cent behind.
- 5 per cent of annual forage are reported as ahead, 89 per cent as normal and 6 per cent behind.

Most crops are still developing and are not ready to harvest. However, a few cereal and early seeded pulse crops are reaching maturity and starting to be harvested.

- 2 per cent of spring wheat;
- 4 per cent of oats; and
- 2 per cent of peas have been harvested.

While harvest has begun, less than 1 per cent of crops in the southeast have been harvested.

Livestock producers are noting that most pastures are in fair to good condition. Currently, 1 per cent of pastures are in excellent condition, 51 per cent are good, 32 per cent are fair, 14 per cent are poor, and 2 per cent are in very poor condition.

Minor to moderate crop damage was reported in the region from hot temperatures and dry conditions, and gophers. Insect damage from aphids, bertha armyworms and grasshoppers were also reported in certain areas in the region.

Southwestern Saskatchewan:

- Census Division 3 – Assiniboia, Gravelbourg, Mankota, Ponteix and Rockglen areas
- Census Division 4 – Cadillac, Consul, Eastend, Maple Creek and Val Marie areas
- Census Division 7 – Beechy, Central Butte, Craik, Herbert, Hodgeville and Moose Jaw areas
- Census Division 8 – Cabri, Elrose, Fox Valley, Leader, Swift Current and Tompkins areas

Much of the region, like most areas across the province, experienced hot, dry weather conditions. A few areas received scattered precipitation, with most areas receiving none. The

highest rainfall in the region was recorded in the Reno area, which was 12 millimeters (mm). The Lake Johnston and Arlington areas followed closely with 11 mm and 10 mm, respectively.

The lack of precipitation significantly reduced topsoil moisture levels, with topsoil moisture mostly rated as short in much of the region.

Cropland topsoil moisture is:

- 23 per cent adequate;
- 44 per cent short; and
- 33 per cent very short.

Hayland topsoil moisture is:

- 18 per cent adequate;
- 56 per cent short; and
- 26 per cent very short.

Pasture topsoil moisture is:

- 15 per cent adequate;
- 60 per cent short; and
- 25 per cent very short.

Southwest Saskatchewan Crop Development			
Crop	% Ahead	% Normal	% Behind
Fall Cereals	5%	90%	5%
Spring Cereals	11%	84%	5%
Oilseeds	12%	80%	8%
Pulse Crops	14%	81%	5%
Perennial Forage	1%	99%	0%
Annual Forage	1%	98%	1%

Hot and dry weather continues to advance crop development in the region. While most crops are rated at normal development, a significant proportion is reported to be ahead of normal development for this time of year.

Crop development for this time of year in the region is rated as follows:

- 5 per cent of fall cereals are reported as ahead, 90 per cent as normal and 5 per cent behind.
- 11 per cent of spring cereals are reported as ahead, 84 per cent as normal and 5 per cent behind.
- 12 per cent of oilseeds are reported as ahead, 80 per cent as normal and 8 per cent behind.
- 14 per cent of pulse crops are reported as ahead, 81 per cent as normal and 5 per cent behind.

- 1 per cent of perennial forage are reported as ahead and 99 per cent as normal.
- 1 per cent of annual forage are reported as ahead, 98 per cent as normal and 1 per cent behind.

Winter cereal harvest is in full swing as large portions of winter wheat and fall rye have either been harvested or are ready to harvest. In total, 5 per cent of crops in the southwest have been harvested, making the southwest region the most advanced to date.

- Winter cereals harvested:
 - Winter wheat: 22 percent; and
 - Fall rye: 36 percent.
- Spring cereals harvested:
 - Triticale: 45 per cent;
 - Barley: 2 per cent; and
 - Oats: 26 per cent.

Pasture conditions in the region are mostly rated as fair to good. Currently, 3 per cent of pastures are in excellent condition, 39 per cent are good, 44 per cent are fair, 12 per cent are poor, and 2 per cent are in very poor conditions.

Hot temperatures and dry conditions were the major cause of crop damage in the region, with some areas reporting moderate to severe damage. Minor crop damage from wind was also reported in certain areas.

East-Central Saskatchewan:

- Census Division 9 – Calder, Canora, Pelly, Preeceville, Sheho and Yorkton areas
- Census Division 10 – Foam Lake, Kelliher, Leroy, Raymore and Wadena areas
- Census Division 11 – Davidson, Colonsay, Langham, Lanigan, Nokomis, Outlook and Saskatoon areas

The region, like most areas in the province, experienced hot, dry weather conditions which have accelerated crop development. Only a few areas in the east-central region recorded some precipitation, with the Invermay area recording 20 mm. This was followed by the Calder and Good Lake areas receiving only 4 mm. Aside from a few other areas which received between 1 to 3 mm of rain, no rainfall was recorded in the rest of the region.

Despite the hot, dry weather conditions, topsoil moisture levels are mostly rated as adequate.

Cropland topsoil moisture is:

- 8 per cent surplus;
- 63 per cent adequate;
- 26 per cent short; and
- 3 per cent very short.

Hayland topsoil moisture is:

- 6 per cent surplus;
- 73 per cent adequate;
- 11 per cent short; and
- 10 per cent very short.

Pasture topsoil moisture is:

- 7 per cent surplus;
- 70 per cent adequate;
- 13 per cent short; and
- 10 per cent very short.

East-Central Saskatchewan Crop Development			
Crop	% Ahead	% Normal	% Behind
Fall Cereals	1%	85%	14%
Spring Cereals	4%	59%	37%
Oilseeds	3%	58%	39%
Pulse Crops	10%	78%	12%
Perennial Forage	4%	72%	24%
Annual Forage	1%	84%	15%

Most of the crops in the region are reported to be at normal stage, with a fair proportion, particularly oilseeds and spring cereals reported as behind for this time of year.

Crop development for this time of year in the region is rated as follows:

- 1 per cent of fall cereals are reported as ahead, 85 per cent as normal and 14 per cent behind.
- 4 per cent of spring cereals are reported as ahead, 59 per cent as normal and 37 per cent behind.
- 3 per cent of oilseeds are reported as ahead, 58 per cent as normal and 39 per cent behind.
- 10 per cent of pulse crops are reported as ahead, 78 per cent as normal and 12 per cent behind.
- 4 per cent of perennial forage are reported as ahead, 72 per cent as normal and 24 per cent behind.
- 1 per cent of annual forage is reported as ahead, 84 per cent as normal and 15 per cent behind.

Most of the spring-seeded crops are not yet ready to harvest; however, fall cereals are starting to reach maturity. Currently, only 15 per cent of fall rye has been harvested in the

region. With hot weather advancing crop maturity, producers are preparing equipment for harvest in the upcoming weeks.

Pasture conditions are mostly fair to good, with a notable proportion rated as excellent. Currently, 17 per cent of pastures are in excellent condition, 56 per cent are good, 15 per cent are fair, 9 per cent are poor, and 3 per cent are very poor. Rain is needed in drier areas to improve subpar pasture conditions.

Minor to moderate crop damage was reported in the region from hot temperatures and dry conditions, with some areas reporting severe damage. Minor damage from wind and insects such as cabbage seedpod weevils and bertha armyworms were also reported. Some notable damage from rust disease was also reported.

West-Central Saskatchewan:

- Census Division 12 – Biggar, Delisle, Rosetown and Sonningdale areas
- Census Division 13 – Cut Knife, Kerrobert, Kindersley, Macklin, Plenty and Wilkie areas

Scattered showers were received in a few areas in the region with most areas receiving none. The Milden area received the most rain with 31 mm, followed by the Manitou Lake area with 12 mm and 10 mm in the Hillsdale area.

The lack of significant rainfall impacted topsoil moisture levels. Topsoil moisture in the region is mostly rated as adequate, with a notable proportion rated as short.

Cropland topsoil moisture is:

- 57 per cent adequate;
- 42 per cent short; and
- 1 per cent very short.

Hayland topsoil moisture is:

- 54 per cent adequate;
- 43 per cent short; and
- 3 per cent very short.

Pasture topsoil moisture is:

- 1 per cent surplus;
- 49 per cent adequate;
- 47 per cent short; and
- 3 per cent very short.

West-Central Saskatchewan Crop Development			
Crop	% Ahead	% Normal	% Behind
Fall Cereals	0%	83%	17%
Spring Cereals	9%	74%	17%
Oilseeds	6%	77%	17%
Pulse Crops	5%	79%	16%
Perennial Forage	0%	75%	25%
Annual Forage	0%	74%	26%

Most of the crops in the west-central are at normal growth stages with a notable proportion behind their development stage for this time of year.

Crop development for this time of year in the region is rated as follows:

- 83 per cent of fall cereals are reported as normal and 17 per cent behind.
- 9 per cent of spring cereals are reported as ahead, 74 per cent as normal and 17 per cent behind.
- 6 per cent of oilseeds are reported as ahead, 77 per cent as normal and 17 per cent behind.
- 5 per cent of pulse crops are reported as ahead, 79 per cent as normal and 16 per cent behind.
- 75 per cent of perennial forage are reported as normal and 25 per cent behind.
- 74 per cent of annual forage are reported as normal and 26 per cent behind.

Most crops are still developing and are not ready to be harvested. Only 1 per cent of oats and barley are reported as harvested in the region. With the hot weather helping to advance crop maturity in the region, producers are looking forward to some much-needed rain in the coming days to help with grain filling for the later seeded crops. Producers are also getting equipment ready for harvest.

Pasture conditions in west-central are mostly rated as fair to good. Currently, 9 per cent of pastures are in excellent condition, 59 per cent of pastures are in good condition, 31 per cent are fair, and 1 per cent are in poor condition.

Crop damage in the region was mostly caused by hot temperatures, dry conditions, wind and gophers. Minor to moderate crop damage from insects such as diamondback and bertha armyworms were also reported in the region.

Northeastern Saskatchewan:

- Census Division 14 – Choiceland, Hudson Bay, Kelvington, Melfort, and Nipawin areas
- Census Division 15 – Cudworth, Humboldt, Kinistino, Prince Albert, Rosthern and St. Brieux areas

Similar to most areas across the province, hot, dry conditions were experienced in the region, with a few areas receiving scattered showers. The 397A area received the highest precipitation of 12 mm, followed closely by the 428 area with 11 mm. The 459 and 463 areas received 8 mm and 6 mm of rain, respectively, taking a toll on many crops in the northeast, and rain is needed soon to support seed development. Producers are preparing for harvest as some expect to begin in the next couple of weeks.

Despite the hot, dry conditions, topsoil moisture levels remain adequate with a few rated as surplus.

Cropland topsoil moisture is:

- 2 per cent surplus;
- 75 per cent adequate; and
- 23 per cent short.

Hayland topsoil moisture is:

- 1 per cent surplus;
- 74 per cent adequate; and
- 25 per cent short.

Pasture topsoil moisture is:

- 5 per cent surplus;
- 70 per cent adequate; and
- 25 per cent short.

Northeast Saskatchewan Crop Development			
Crop	% Ahead	% Normal	% Behind
Fall Cereals	0%	75%	25%
Spring Cereals	1%	69%	30%
Oilseeds	2%	62%	36%
Pulse Crops	3%	72%	25%
Perennial Forage	0%	80%	20%
Annual Forage	0%	82%	18%

The warm weather has helped advance crop development in the region, with most of the crops reported to be at their normal stage. A notable proportion of crops are also reported to be behind in their development for this time of year.

Crop development for this time of year in the region is rated as follows:

- 75 per cent of fall cereals are reported as normal and 25 per cent behind.
- 1 per cent of spring cereals are reported as ahead, 69 per cent as normal and 30 per cent behind.

- 2 per cent of oilseeds are reported as ahead, 62 per cent as normal and 36 per cent behind.
- 3 per cent of pulse crops are reported as ahead, 72 per cent as normal and 25 per cent behind.
- 80 per cent of perennial forage are reported as normal and 20 per cent behind.
- 82 per cent of annual forage are reported as normal and 18 per cent behind.

Although hot, dry conditions have helped accelerate crop maturity in the region and producers are getting ready for harvest, it will be a while until most crops in the region are ready for harvest. So far, only 1 per cent of barley has been harvested in the region.

Pasture conditions in the region are mostly rated as good. While 5 per cent of pastures are in excellent condition, 72 per cent of pastures are reported to be in good condition, with 23 per cent in fair condition.

Like in most parts of the province, hot, dry weather conditions were reported as the major cause of crop damage in the northeast region. Minor crop damage from aphids was also reported.

Northwestern Saskatchewan:

- Census Division 16 – Blaine Lake, Canwood, North Battleford, Radisson and Spiritwood areas
- Census Division 17 – Glaslyn, Maidstone, Meadow Lake, Pierceland and St. Walburg areas

Unlike other regions across the province, several areas in the northwest received at least 10 mm of rain. The Beaver River area received the most with 28 mm, followed by the Meadow Lake area with 18 mm. The Eldon and Meeting Lake areas also received 14 mm and 12 mm of rain respectively.

Topsoil moisture in the region is mostly rated adequate.

Cropland topsoil moisture is:

- 2 per cent surplus;
- 95 per cent adequate; and
- 3 per cent short.

Hayland topsoil moisture is:

- 2 per cent surplus;
- 95 per cent adequate; and
- 3 per cent short.

Pasture topsoil moisture is:

- 1 per cent surplus;

- 95 per cent adequate; and
- 4 per cent short.

Northwest Saskatchewan Crop Development			
Crop	% Ahead	% Normal	% Behind
Fall Cereals	0%	25%	75%
Spring Cereals	4%	64%	32%
Oilseeds	4%	57%	39%
Pulse Crops	6%	68%	26%
Perennial Forage	0%	82%	18%
Annual Forage	2%	72%	26%

While the majority of the crops are reported to be at their normal stage of development, a substantial amount, particularly in the fall cereals, are reported to be behind their development for this time of year.

Crop development for this time of year in the region is rated as follows:

- 25 per cent of fall cereals are reported as normal and 75 per cent behind.
- 4 per cent of spring cereals are reported as ahead, 64 per cent as normal and 32 per cent behind.
- 4 per cent of oilseeds are reported as ahead, 57 per cent as normal and 39 per cent behind.
- 6 per cent of pulse crops are reported as ahead, 68 per cent as normal and 26 per cent behind.
- 82 per cent of perennial forage are reported as normal and 18 per cent behind.
- 2 per cent of annual forage are reported as ahead, 72 per cent as normal and 26 per cent behind.

Most crops are still developing and are not ready to be harvested. The recent hot, dry weather conditions will be helpful in accelerating crop development and maturity in the region.

Pasture conditions are mostly fair to good, with a notable proportion reported to be in excellent condition. 11 per cent of pastures are reported to be in excellent condition, 70 per cent are in good condition, 18 per cent are fair, and 1 per cent are in poor condition.

Major crop damage in the region was caused by hot weather, wind and gopher. Bertha armyworms were also reported to cause minor crop damage in some areas in the region.

Saskatchewan Crop Development (for the period of July 28 to August 3, 2026)

Provincial Crop Development

Crop	% Ahead	% Normal	% Behind
Fall Cereals	4%	86%	10%
Spring Cereals	7%	71%	22%
Oilseeds	6%	65%	29%
Pulse Crops	10%	77%	13%
Perennial Forage	2%	83%	15%
Annual Forage	1%	78%	21%

Southeast Saskatchewan Crop Development

Crop	% Ahead	% Normal	% Behind
Fall Cereals	6%	89%	5%
Spring Cereals	10%	76%	14%
Oilseeds	8%	74%	18%
Pulse Crops	10%	78%	12%
Perennial Forage	4%	90%	6%
Annual Forage	5%	89%	6%

Southwest Saskatchewan Crop Development

Crop	% Ahead	% Normal	% Behind
Fall Cereals	5%	90%	5%
Spring Cereals	11%	84%	5%
Oilseeds	12%	80%	8%
Pulse Crops	14%	81%	5%
Perennial Forage	1%	99%	0%
Annual Forage	1%	98%	1%

East-Central Saskatchewan Crop Development

Crop	% Ahead	% Normal	% Behind
Fall Cereals	1%	85%	14%
Spring Cereals	4%	59%	37%
Oilseeds	3%	58%	39%
Pulse Crops	10%	78%	12%
Perennial Forage	4%	72%	24%
Annual Forage	1%	84%	15%

West-Central Saskatchewan Crop Development

Crop	% Ahead	% Normal	% Behind
Fall Cereals	0%	83%	17%
Spring Cereals	9%	74%	17%
Oilseeds	6%	77%	17%
Pulse Crops	5%	79%	16%
Perennial Forage	0%	75%	25%
Annual Forage	0%	74%	26%

Northeast Saskatchewan Crop Development

Crop	% Ahead	% Normal	% Behind
Fall Cereals	0%	75%	25%
Spring Cereals	1%	69%	30%
Oilseeds	2%	62%	36%
Pulse Crops	3%	72%	25%
Perennial Forage	0%	80%	20%
Annual Forage	0%	82%	18%


Northwest Saskatchewan Crop Development

Crop	% Ahead	% Normal	% Behind
Fall Cereals	0%	25%	75%
Spring Cereals	4%	64%	32%
Oilseeds	4%	57%	39%
Pulse Crops	6%	68%	26%
Perennial Forage	0%	82%	18%
Annual Forage	2%	72%	26%

Provincial Harvest Progress	
Per cent Harvested for All Crops	
for the Period up to August 3, 2026	
Region	Harvest Completed
Southeast	0%
Southwest	5%
East-central	0%
West-central	0%
Northeast	0%
Northwest	0%
Provincial	2%

Weekly Rainfall

from July 28 to August 3, 2026



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Weekly Rainfall Summary

(reported in millimeters)

1 inch=25 mm

for the period from July 28 to August 3, 2026

Census Division	RM No.	RM Name	Past Week	Since 1-Apr	Census Division	RM No.	RM Name	Past Week	Since 1-Apr	Census Division	RM No.	RM Name	Past Week	Since 1-Apr
1	3	ENNISKILLEN	0	287	8	138	WEBB	N/A	224	14	366	KELVINGTON	0	241
1	4	COALFIELDS	0	378	8	139	GULL LAKE	3	208	14	367	PONASS LAKE	N/A	306
1	32	RECIPROCITY	7	278	8	168	RIVERSIDE	0	193	14	394	HUDSON BAY	0	222
1	34	BROWNING	N/A	290	8	228	LACADENA	1	233	14	395	PORCUPINE	N/A	273
1	94	HAZELWOOD	N/A	0	8	229	MIRY CREEK	N/A	152	14	397	BARRIER VALLEY	0	184
1	95	GOLDEN WEST	N/A	225	8	231	HAPPYLAND	2	161	14	428	STAR CITY	11	222
1	2A	MOUNT PLEASANT	5	200	8	259	SNIFE LAKE	0	189	14	456	ARBORFIELD	0	257
2	10	HAPPY VALLEY	0	196	8	138A	WEBB	3	154	14	457	CONNAUGHT	0	67
2	38	LAURIER	0	248	8	228A	LACADENA	0	64	14	486	MOOSE RANGE	0	212
2	66	GRIFFIN	N/A	294	8	257A	MONET	0	188	14	487	NIPAWIN	N/A	225
2	67	WEYBURN	N/A	101	8	259A	SNIFE LAKE	N/A	141	14	488	TORCH RIVER	0	226
2	70	KEY WEST	N/A	12	9	241	CALDER	3	383	14	394A	HUDSON BAY	1	245
2	100	ELMSTHORPE	0	209	9	243	WALLACE	0	232	14	397A	BARRIER VALLEY	12	235
2	100A	ELMSTHORPE	2	223	9	245	GARRY	N/A	158	14	488A	TORCH RIVER	N/A	116
2	38A	LAURIER	0	215	9	273	SLIDING HILLS	0	318	14	488B	TORCH RIVER	N/A	5
3	73	STONEHENGE	0	209	9	274	GOOD LAKE	4	244	15	369	ST. PETER	1	208
3	74	WOOD RIVER	0	207	9	301	ST. PHILIPS	0	143	15	371	BAYNE	0	246
3	75	PINTO CREEK	0	145	9	305	INVERMAY	20	310	15	372	GRANT	0	182
3	76	AUVERGNE	1	199	9	331	LIVINGSTON	3	238	15	373	ABERDEEN	0	186
3	101	TERRELL	N/A	0	9	241A	CALDER	4	470	15	399	LAKE LENORE	3	235
3	102	LAKE JOHNSTON	11	95	9	245A	GARRY	N/A	211	15	400	THREE LAKES	0	311
3	106	WHISKA CREEK	0	178	10	246	ITUNA BON ACCORD	0	131	15	402	FISH CREEK	0	171
3	74A	KEY WEST	0	170	10	247	KELLROSS	2	216	15	429	FLETT'S SPRINGS	N/A	214
4	51	RENO	12	130	10	248	TOUCHWOOD	0	203	15	430	INVERGORDON	0	120
4	79	ARLINGTON	0	218	10	277	EMERALD	0	206	15	459	KINISTINO	8	259
4	110	PIAPOT	0	217	10	279	MOUNT HOPE	0	189	15	461	PRINCE ALBERT	N/A	224
4	77A	WISE CREEK	0	152	10	307	ELFROS	2	399	15	463	DUCK LAKE	6	236
4	78A	GRASSY CREEK	N/A	127	10	309	PRAIRIE ROSE	2	255	15	491	BUCKLAND	0	105
4	79A	ARLINGTON	10	236	10	336	SASMAN	N/A	299	15	520	PADDOCKWOOD	0	20
5	122	MARTIN	0	257	10	339	LEROY	1	236	15	521	LAKELAND	0	20
5	124	KINGSLEY	7	251	10	246A	ITUNA BON ACCORD	1	105	15	371A	BAYNE	0	176
5	151	ROCANVILLE	1	239	10	248A	TOUCHWOOD	0	345	15	371B	BAYNE	0	233
5	155	WOLSELEY	2	196	10	276A	FOAM LAKE	0	341	15	399A	LAKE LENORE	N/A	16
5	181	LANGENBURG	5	423	10	276B	FOAM LAKE	0	454	15	403A	ROSTHERN	5	249
5	183	FERTILE BELT	N/A	273	10	276C	FOAM LAKE	0	355	15	403C	ROSTHERN	0	245
5	211	CHURCHBRIDGE	0	296	10	276D	FOAM LAKE	0	260	15	461A	PRINCE ALBERT	N/A	192
5	213	SALTCOATS	1	329	10	277A	EMERALD	N/A	270	15	461B	PRINCE ALBERT	N/A	2
5	215	STANLEY	0	218	10	279A	MOUNT HOPE	2	158	16	434	BLAINE LAKE	N/A	147
5	125A	CHESTERFIELD	0	188	11	251	BIG ARM	N/A	232	16	435	REDBERRY	5	295
5	154A	ELCAPO	2	213	11	280	WREFOED	N/A	83	16	436	DOUGLAS	0	192
5	183A	FERTILE BELT	0	332	11	283	ROSEDALE	0	255	16	437	NORTH BATTLEFORD	5	337
5	211A	CHURCHBRIDGE	0	317	11	284	RUDY	0	65	16	466	MEETING LAKE	12	294
6	127	FRANCIS	0	218	11	314	DUNDURN	0	201	16	467	ROUND HILL	10	254
6	128	LAJORD	0	111	11	344	CORMAN PARK	N/A	154	16	493	SHELLBROOK	N/A	174
6	130	REDBURN	0	238	11	282A	McCRANEY	0	155	16	494	CANWOOD	3	116
6	156	INDIAN HEAD	9	243	12	286	MILDEN	31	205	16	497	MEDSTEAD	6	267
6	160	PENSE	0	160	12	287	ST. ANDREWS	0	160	16	437A	NORTH BATTLEFORD	6	279
6	186	ABERNETHY	0	231	12	288	PLEASANT VALLEY	0	155	16	467A	ROUND HILL	8	307
6	190	DUFFERIN	N/A	178	12	316	HARRIS	0	133	17	468	MEOTA	10	215
6	216	TULLYMET	N/A	10	12	317	MARRIOTT	0	203	17	470	PAYNTON	N/A	76
6	217	LIPTON	0	329	12	345	VANSCHOY	0	130	17	471	ELDON	14	274
6	219	LONGLAKETON	0	238	12	346	PERDUE	0	176	17	498	PARKDALE	8	310
6	221	SARNIA	1	213	12	347	BIGGAR	0	239	17	499	MERVIN	N/A	223
6	127B	FRANCIS	N/A	116	12	376	EAGLE CREEK	0	225	17	502	BRITANNIA	7	191
6	159A	SHERWOOD	N/A	131	12	377	GLENSIDE	2	207	17	561	LOON LAKE	6	163
6	159B	SHERWOOD	1	189	12	285A	FERTILE VALLEY	0	144	17	588	MEADOW LAKE	18	318
6	160A	PENSE	0	139	13	290	KINDERSLEY	N/A	166	17	622	BEAVER RIVER	28	265
6	190A	DUFFERIN	0	120	13	292	MILTON	3	333	17	498A	PARKDALE	10	262
6	190B	DUFFERIN	N/A	204	13	321	PRAIRIEDALE	N/A	242	17	501A	FRENCHMAN BUTTE	9	239
6	190C	DUFFERIN	0	246	13	350	MARIPOSA	0	200	17	501C	FRENCHMAN BUTTE	N/A	50
6	216A	TULLYMET	N/A	75	13	351	PROGRESS	1	229	17	561A	LOON LAKE	5	174
6	217A	LIPTON	N/A	2	13	379	REFORD	N/A	93					
6	219A	LONGLAKETON	0	226	13	382	EYE HILL	3	217					
6	219B	LONGLAKETON	N/A	60	13	409	BUFFALO	0	270					
6	220A	McKILLOP	0	161	13	440	HILLSDALE	10	241					
6	220B	McKILLOP	0	0	13	442	MANITOU LAKE	12	232					
7	132	HILLSBOROUGH	0	299	13	292A	MILTON	0	171					
7	136	COULEE	N/A	151	13	320A	OAKDALE	0	167					
7	161	MOOSE JAW	3	142	13	321A	PRAIRIEDALE	N/A	35					
7	162	CARON	0	210	13	409A	BUFFALO	0	227					
7	165	MORSE	0	250										
7	193	EYEBROW	0	242										
7	223	HURON	1	185										
7	132A	HILLSBOROUGH	0	118										
7	136A	COULEE	N/A	0										
7	161A	MOOSE JAW	N/A	38										
7	162A	CARON	N/A	183										
7	222A	CRAIK	3	228										
7	223A	HURON	3	177										

Municipality No: A, B, C and D - more than one reporter


These precipitation amounts represent point locations within each municipality and do not necessarily reflect the whole R. M.
N/A indicates that rainfall was not reported for the week

Cumulative Rainfall

from April 1 to August 3, 2026

Rainfall (mm)


0.0 - 25.0	100.1 - 125.0	200.1 - 225.0	300.1 - 325.0	400.1 - 425.0
25.1 - 50.0	125.1 - 150.0	225.1 - 250.0	325.1 - 350.0	425.1 - 450.0
50.1 - 75.0	150.1 - 175.0	250.1 - 275.0	350.1 - 375.0	450.1 - 475.0
75.1 - 100.0	175.1 - 200.0	275.1 - 300.0	375.1 - 400.0	475.1 +



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Cropland Topsoil Moisture Conditions


from July 28 to August 3, 2026



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Hay Topsoil Moisture Conditions


from July 28 to August 3, 2026



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Pasture Topsoil Moisture Conditions


from July 28 to August 3, 2026



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Pasture Conditions


from July 28 to August 3, 2026



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.


Maximum Temperature

from July 28 to August 3, 2026



Minimum Temperature

from July 28 to August 3, 2026



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.