

Crop Report

For the Period August 4 to August 10, 2026

Published by the Ministry of Agriculture

ISSN 0701 7085

Report number 15, August 13, 2026

Crops continue to mature across the province. Cooler weather slowed harvest progress, which remains at 2 per cent complete, and crop development in some areas; however, it also helped maintain crop condition and reduce stress in areas that experienced hot, dry conditions over the past few weeks. Scattered showers provided much-needed moisture, but overall conditions remain dry. Additional precipitation would be beneficial, particularly for later-seeded crops as they continue to progress through the grain-filling stage.

Variable rainfall was received across the province. While a few areas received significant precipitation, most regions recorded little to no rainfall. The highest amount was reported in the Chesterfield and Milton areas, where 20 millimetres (mm) of rain fell, followed by the Milden area with 16 mm and the Harris, Arborfield, Nipawin, Round Hill and Hillsdale areas at 15 mm.

The localized showers improved topsoil moisture conditions in some areas; however, a significant proportion of regions across the province rated topsoil moisture levels as short.

Cropland topsoil moisture is:

- 1 per cent surplus;
- 48 per cent adequate;
- 43 per cent short; and
- 8 per cent very short.

Hayland topsoil moisture is:

- 1 per cent surplus;
- 49 per cent adequate;
- 42 per cent short; and
- 8 per cent very short.

One year ago

Widespread rainfall was received across the province. While the precipitation was timely and beneficial for later-seeded crops, it also contributed to delayed crop development and slowed harvest progress in many areas. As a result, only one per cent of crops had been harvested, primarily winter cereals and pulse crops. Haying and silaging operations were largely complete or nearing completion across the province.

Follow the 2026 Crop Report on X
@SKAgriculture

Harvest Progress in SK

Per Cent Combined
Historical All Crops

Aug 10/26	2
Aug 11/25	1
Aug 12/24	6
Aug 14/23	9
Aug 15/22	5
Aug 9/21	7
5 year avg. (2021-2025)	6
10 year avg. (2016-2025)	5

For further information, contact Augustine Osei, PhD, PAg,
Crops Extension Specialist, Extension Services Branch,
Toll Free: 1-866-457-2377 or 306-520-5761, Email: cropreport@gov.sk.ca.
Also available on the Ministry of Agriculture website at saskatchewan.ca/crop-report.



**Agriculture and
Agri-Food Canada**

SCIC
SASKATCHEWAN CROP
INSURANCE CORPORATION

Saskatchewan

Crop Report

For the Period August 4 to August 10, 2026

Published by the Ministry of Agriculture

ISSN 0701 7085

Report number 15, August 13, 2026

Pasture topsoil moisture is:

- 1 per cent surplus;
- 39 per cent adequate;
- 46 per cent short; and
- 14 per cent very short.

Cool weather conditions slowed harvest operations in areas where harvest has begun. In most regions, however, crops continue to develop and are not yet ready for harvest. Provincial harvest progress remains at 2 per cent complete, largely limited to cereal and early-seeded pulse crops. This is below both the five-year average of 6 per cent and ten-year average of 5 per cent. However, harvest progress is ahead of last year, when only one per cent of the crop had been harvested. Desiccation operations are underway in pulse crops, while many crops across the province continue to mature.

- Winter cereals harvested:
 - Winter wheat: 18 per cent; and
 - Fall rye: 27 per cent.
- Spring cereals harvested:
 - Triticale: 37 per cent;
 - Barley: 4 per cent;
 - Spring wheat: 1 per cent;
 - Oats: 2 per cent; and
 - Canary seed: 1 per cent.
- Pulses harvested:
 - Lentil: 2 per cent; and
 - Field peas: 5 per cent.

Most producers have either completed or are wrapping up final haying and silage operations. Provincially, dryland alfalfa yields are averaging 1.90 tonnes (t) per acre, while greenfeed yields are averaging 2.79 t per acre. Under irrigated conditions, alfalfa yields are averaging 3.90 t per acre and greenfeed yields are averaging 6.85 t per acre. Provincial silage yields are currently estimated at 7.00 t per acre.

For further information, contact Augustine Osei, PhD, PAg,
Crops Extension Specialist, Extension Services Branch,
Toll Free: 1-866-457-2377 or 306-520-5761, Email: cropreport@gov.sk.ca.
Also available on the Ministry of Agriculture website at saskatchewan.ca/crop-report.



**Agriculture and
Agri-Food Canada**

SCIC
SASKATCHEWAN CROP
INSURANCE CORPORATION

Saskatchewan

Crop Report

For the Period August 4 to August 10, 2026

Published by the Ministry of Agriculture

ISSN 0701 7085

Report number 15, August 13, 2026

Crop damage was reported in areas experiencing hot, dry conditions, with some regions indicating minor to moderate crop stress. There were also reports of bertha armyworms, wind, gophers and sclerotinia disease causing minor crop damage in certain areas.

Harvest is a busy time across Saskatchewan and safety should remain a top priority for everyone. Producers are reminded to be aware of potential hazards, including fire risks and overhead powerlines, and to take appropriate preventative measures to reduce the likelihood of incidents. As harvest activity continues to increase, there will be more farm machinery travelling on public roadways. Motorists are encouraged to exercise extra patience and caution when approaching or passing large agricultural equipment.

For any crop or livestock questions, producers are encouraged to call the Agriculture Knowledge Centre, toll free at 1-866-457-2377.

This can also be a stressful time of year for producers, particularly as weather conditions remain unpredictable and harvest progress is impacted. Support is available through the Farm Stress Line at 1-800-667-4442, which provides confidential assistance to producers and their families.

For further information, contact Augustine Osei, PhD, PAg,
Crops Extension Specialist, Extension Services Branch,
Toll Free: 1-866-457-2377 or 306-520-5761, Email: cropreport@gov.sk.ca.
Also available on the Ministry of Agriculture website at saskatchewan.ca/crop-report.



**Agriculture and
Agri-Food Canada**



SCIC
SASKATCHEWAN CROP
INSURANCE CORPORATION

Saskatchewan

Harvest Progress

from April 1 to August 10, 2026

Harvest Progress



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Southeast Saskatchewan	
Census Division	% Harvested (August 10, 2026)
1	2%
2	3%
5	0%
6	1%
Region average	1%

Southwest Saskatchewan	
Census Division	% Harvested (August 10, 2026)
3	2%
4	6%
7	9%
8	3%
Region average	5%

East-Central Saskatchewan	
Census Division	% Harvested (August 10, 2026)
9	0%
10	0%
11	2%
Region average	1%

West-Central Saskatchewan	
Census Division	% Harvested (August 10, 2026)
12	0%
13	0%
Region average	0%

Northeast Saskatchewan	
Census Division	% Harvested (August 10, 2026)
14	0%
15	0%
Region average	0%

Northwest Saskatchewan	
Census Division	% Harvested (August 10, 2026)
16	0%
17	0%
Region average	0%

Saskatchewan Harvest Progress - August 10, 2026

*Other - crop that will not be harvested due to weather, insect or disease damage

Provincial								
	Winter Wheat	Fall Rye	Spring Wheat	Durum	Oats	Barley	Flax	Canola
Standing	53%	42%	99%	99%	97%	95%	100%	100%
In Swath	0%	11%	0%	0%	0%	0%	0%	0%
Ready To Straight Combine	29%	20%	0%	1%	0%	1%	0%	0%
Combined	15%	20%	1%	0%	1%	2%	0%	0%
Harvested As Feed	3%	7%	0%	0%	2%	2%	0%	0%
Other (Damage/Challenges)	0%	0%	0%	0%	0%	0%	0%	0%
	Triticale	Mustard	Soybean	Lentil	Field Pea	Canaryseed	Chickpea	
Standing	63%	100%	100%	88%	83%	97%	100%	
In Swath	0%	0%	0%	3%	0%	0%	0%	
Ready To Straight Combine	3%	0%	0%	6%	12%	2%	0%	
Combined	0%	0%	0%	2%	5%	1%	0%	
Harvested As Feed	34%	0%	0%	0%	0%	0%	0%	
Other (Damage/Challenges)	0%	0%	0%	1%	0%	0%	0%	
South East								
	Winter Wheat	Fall Rye	Spring Wheat	Durum	Oats	Barley	Flax	Canola
Standing	55%	62%	98%	98%	84%	84%	100%	100%
In Swath	0%	0%	0%	0%	0%	0%	0%	0%
Ready To Straight Combine	44%	19%	0%	1%	2%	4%	0%	0%
Combined	1%	3%	2%	0%	12%	10%	0%	0%
Harvested As Feed	0%	16%	0%	0%	2%	2%	0%	0%
Other (Damage/Challenges)	0%	0%	0%	1%	0%	0%	0%	0%
	Triticale	Mustard	Soybean	Lentil	Field Pea	Canaryseed	Chickpea	
Standing	100%	100%	100%	81%	64%	96%	100%	
In Swath	0%	0%	0%	0%	0%	0%	0%	
Ready To Straight Combine	0%	0%	0%	12%	23%	2%	0%	
Combined	0%	0%	0%	2%	12%	2%	0%	
Harvested As Feed	0%	0%	0%	0%	0%	0%	0%	
Other (Damage/Challenges)	0%	0%	0%	5%	1%	0%	0%	
South West								
	Winter Wheat	Fall Rye	Spring Wheat	Durum	Oats	Barley	Flax	Canola
Standing	16%	31%	99%	97%	68%	92%	100%	100%
In Swath	0%	17%	0%	0%	3%	0%	0%	0%
Ready To Straight Combine	31%	23%	1%	3%	3%	2%	0%	0%
Combined	45%	19%	0%	0%	0%	1%	0%	0%
Harvested As Feed	8%	10%	0%	0%	26%	5%	0%	0%
Other (Damage/Challenges)	0%	0%	0%	0%	0%	0%	0%	0%
	Triticale	Mustard	Soybean	Lentil	Field Pea	Canaryseed	Chickpea	
Standing	40%	100%	100%	83%	67%	100%	100%	
In Swath	0%	0%	0%	4%	0%	0%	0%	
Ready To Straight Combine	3%	0%	0%	10%	23%	0%	0%	
Combined	0%	0%	0%	3%	10%	0%	0%	
Harvested As Feed	57%	0%	0%	0%	0%	0%	0%	
Other (Damage/Challenges)	0%	0%	0%	0%	0%	0%	0%	

Saskatchewan Harvest Progress - August 10, 2026

*Other - crop that will not be harvested due to weather, insect or disease damage

East-Central								
	Winter Wheat	Fall Rye	Spring Wheat	Durum	Oats	Barley	Flax	Canola
Standing	64%	5%	100%	100%	100%	98%	100%	100%
In Swath	0%	2%	0%	0%	0%	0%	0%	0%
Ready To Straight Combine	36%	1%	0%	0%	0%	1%	0%	0%
Combined	0%	92%	0%	0%	0%	0%	0%	0%
Harvested As Feed	0%	0%	0%	0%	0%	1%	0%	0%
Other (Damage/Challenges)	0%	0%	0%	0%	0%	0%	0%	0%
	Triticale	Mustard	Soybean	Lentil	Field Pea	Canaryseed	Chickpea	
Standing	95%	100%	100%	86%	95%	100%	100%	
In Swath	0%	0%	0%	0%	1%	0%	0%	
Ready To Straight Combine	0%	0%	0%	12%	3%	0%	0%	
Combined	5%	0%	0%	2%	1%	0%	0%	
Harvested As Feed	0%	0%	0%	0%	0%	0%	0%	
Other (Damage/Challenges)	0%	0%	0%	0%	0%	0%	0%	
West-Central								
	Winter Wheat	Fall Rye	Spring Wheat	Durum	Oats	Barley	Flax	Canola
Standing	67%	78%	100%	100%	97%	92%	100%	99%
In Swath	2%	2%	0%	0%	0%	0%	0%	1%
Ready To Straight Combine	22%	13%	0%	0%	0%	0%	0%	0%
Combined	10%	7%	0%	0%	0%	6%	0%	0%
Harvested As Feed	0%	0%	0%	0%	3%	2%	0%	0%
Other (Damage/Challenges)	0%	0%	0%	0%	0%	0%	0%	0%
	Triticale	Mustard	Soybean	Lentil	Field Pea	Canaryseed	Chickpea	
Standing	100%	94%	100%	96%	97%	100%	100%	
In Swath	0%	0%	0%	0%	0%	0%	0%	
Ready To Straight Combine	0%	0%	0%	3%	2%	0%	0%	
Combined	0%	6%	0%	1%	1%	0%	0%	
Harvested As Feed	0%	0%	0%	0%	0%	0%	0%	
Other (Damage/Challenges)	0%	0%	0%	0%	0%	0%	0%	
North East								
	Winter Wheat	Fall Rye	Spring Wheat	Durum	Oats	Barley	Flax	Canola
Standing	66%	66%	100%	100%	100%	99%	100%	100%
In Swath	0%	0%	0%	0%	0%	0%	0%	0%
Ready To Straight Combine	34%	0%	0%	0%	0%	0%	0%	0%
Combined	0%	34%	0%	0%	0%	0%	0%	0%
Harvested As Feed	0%	0%	0%	0%	0%	1%	0%	0%
Other (Damage/Challenges)	0%	0%	0%	0%	0%	0%	0%	0%
	Triticale	Mustard	Soybean	Lentil	Field Pea	Canaryseed	Chickpea	
Standing	100%	100%	100%	99%	100%	100%	100%	
In Swath	0%	0%	0%	1%	0%	0%	0%	
Ready To Straight Combine	0%	0%	0%	0%	0%	0%	0%	
Combined	0%	0%	0%	0%	0%	0%	0%	
Harvested As Feed	0%	0%	0%	0%	0%	0%	0%	
Other (Damage/Challenges)	0%	0%	0%	0%	0%	0%	0%	


Saskatchewan Harvest Progress - August 10, 2026

*Other - crop that will not be harvested due to weather, insect or disease damage

North West								
	Winter Wheat	Fall Rye	Spring Wheat	Durum	Oats	Barley	Flax	Canola
Standing	100%	› Response	100%	Response	100%	100%	100%	100%
In Swath	0%	› Response	0%	Response	0%	0%	0%	0%
Ready To Straight Combine	0%	› Response	0%	Response	0%	0%	0%	0%
Combined	0%	› Response	0%	Response	0%	0%	0%	0%
Harvested As Feed	0%	› Response	0%	Response	0%	0%	0%	0%
Other (Damage/Challenges)	0%	› Response	0%	Response	0%	0%	0%	0%
	Triticale	Mustard	Soybean	Lentil	Field Pea	Canaryseed	Chickpea	
Standing	100%	› Response	No Response(s)	99%	99%	No Response(s)	› Response(s)	
In Swath	0%	› Response	No Response(s)	0%	0%	No Response(s)	› Response(s)	
Ready To Straight Combine	0%	› Response	No Response(s)	0%	0%	No Response(s)	› Response(s)	
Combined	0%	› Response	No Response(s)	0%	0%	No Response(s)	› Response(s)	
Harvested As Feed	0%	› Response	No Response(s)	0%	0%	No Response(s)	› Response(s)	
Other (Damage/Challenges)	0%	› Response	No Response(s)	1%	1%	No Response(s)	› Response(s)	

Saskatchewan Harvest Progress	
Up to August 10, 2026	
Provincial % Combined	
Winter Wheat**	18%
Fall Rye**	27%
Spring Wheat	0%
Durum	0%
Oats**	2%
Barley**	4%
Flax	0%
Canola	0%
Triticale**	34%
Mustard	0%
Soybean	0%
Lentil	2%
Field Pea	5%
Canaryseed	1%
Chickpea	0%
**includes per cent harvested as feed	

Crop Report Regions & Census Divisions



Southeastern Saskatchewan:

- Census Division 1 – Carnduff, Estevan, Lampman, Redvers and Stoughton areas
- Census Division 2 – Avonlea, Fillmore, Minton, Radville and Weyburn areas
- Census Division 5 – Broadview, Esterhazy, Melville and Moosomin areas
- Census Division 6 – Belle Plaine, Cupar, Lumsden, Indian Head, Regina and Rouleau areas

Scattered rainfall across the region slowed harvest and haying operations in some areas. Although the isolated showers and generally cooler temperatures helped maintain crop conditions and reduced stress caused by the hot, dry weather, they also slowed crop development. Additional precipitation will be critical in the coming days, particularly for later-seeded crops.

Apart from the Chesterfield area, which received the highest rainfall amount at 20 millimetres (mm), most locations recorded less than 10 mm of precipitation. The Kingsley and Wolseley areas each received 6 mm. Several parts of the region received little to no rainfall, further impacting topsoil moisture conditions, which are mostly rated as short.

Cropland topsoil moisture is:

- 43 per cent adequate;
- 55 per cent short; and
- 2 per cent very short.

Hayland topsoil moisture is:

- 42 per cent adequate;
- 55 per cent short; and
- 3 per cent very short.

Pasture topsoil moisture is:

- 36 per cent adequate;
- 59 per cent short; and
- 5 per cent very short.

A few crops in the southeast region are reaching maturity, with harvest now underway. Approximately 1 per cent of the crop, primarily cereals and a small number of pulse crops, have been harvested. This is below both the five-year and 10-year average of 6 per cent but is on par with last year, when 1 per cent of the crop had been harvested at this time.

Winter cereals harvested:

- Winter wheat: 1 per cent; and
- Fall rye: 19 per cent.

Spring cereals harvested:

- Barley: 12 per cent;
- Spring wheat: 2 per cent;

- Oats: 14 per cent; and
- Canary seed: 2 per cent.

Pulses harvested:

- Lentil: 2 per cent; and
- Field peas: 12 per cent.

Livestock producers are wrapping up haying and silaging for the season. Average yields for dryland forage crops were reported at 1.89 tonne (t) per acre for alfalfa, 1.71 t per acre for tame hay and 2.73 t per acre for greenfeed. Irrigated forage crops produced higher yields, with both alfalfa and tame hay averaging 6 t per acre, while irrigated greenfeed averaged 9 t per acre. Silage crops yielded an average of 6.88 t per acre across the region.

Minor to severe crop damage was reported in the region from hot, dry conditions, wind and gophers. Minor crop damage from grasshoppers was also reported in certain areas in the region.

Southwestern Saskatchewan:

- Census Division 3 – Assiniboia, Gravelbourg, Mankota, Ponteix and Rockglen areas
- Census Division 4 – Cadillac, Consul, Eastend, Maple Creek and Val Marie areas
- Census Division 7 – Beechy, Central Butte, Craik, Herbert, Hodgeville and Moose Jaw areas
- Census Division 8 – Cabri, Elrose, Fox Valley, Leader, Swift Current and Tompkins areas

The region experienced cool weather and scattered showers in a few isolated areas, while most locations received little to no precipitation. The highest rainfall amount was recorded in the Happyland area, which received only 6 millimetres (mm) of rain. The Caron and Snipe Lake areas followed closely with 5 mm of precipitation.

Limited rainfall across the region significantly reduced topsoil moisture levels, which is now rated as mostly short in much of the region.

Cropland topsoil moisture is:

- 18 per cent adequate;
- 58 per cent short; and
- 24 per cent very short.

Hayland topsoil moisture is:

- 19 per cent adequate;
- 54 per cent short; and
- 27 per cent very short.

Pasture topsoil moisture is:

- 13 per cent adequate;
- 60 per cent short; and
- 27 per cent very short.

Despite being the most advanced region in the province for harvest progress to date, harvest remains at 5 per cent complete, as cool weather conditions continued to slow field operations. This is below both the five-year average of 13 per cent and the 10-year average of 11 per cent. However, it is ahead of last year, when 3 per cent of the crop had been harvested by this time.

Winter cereals harvested:

- Winter wheat: 53 per cent; and
- Fall rye: 36 per cent.

Spring cereals harvested:

- Triticale: 57 per cent;
- Barley: 12 per cent; and
- Oats: 26 per cent.

Pulses harvested:

- Lentil: 3 per cent; and
- Field peas: 10 per cent.

Livestock producers have largely wrapped up haying and silaging this year. Dryland alfalfa yields averaged 1.56 tonne (t) per acre, tame hay yielded 1.30 t per acre and greenfeed yielded 2.35 t per acre on average. Under irrigated conditions, alfalfa yields averaged 2.94 t per acre. Silage crops yielded the highest in the southwest compared to the rest of the province with an average yield of 9.50 t per acre.

Hot, dry conditions were the major cause of crop damage in the region with some areas reporting moderate to severe damage. Minor to moderate crop damage from wind was also reported in certain areas.

East-Central Saskatchewan:

- Census Division 9 – Calder, Canora, Pelly, Preeceville, Sheho and Yorkton areas
- Census Division 10 – Foam Lake, Kelliher, Leroy, Raymore and Wadena areas
- Census Division 11 – Davidson, Colonsay, Langham, Lanigan, Nokomis, Outlook and Saskatoon areas

This region, like most areas in the province, experienced cool weather conditions with some scattered showers in isolated places slowing harvest and haying. Conditions are generally dry and producers are hoping for some much-needed rain to help later seeded crops with grain filling.

The highest amount of precipitation that was received in the region was 9 millimetres (mm) in the Rudy areas. This was followed by the Rosedale areas which received 6 mm and 4 mm in the Calder area.

Despite the lack of rainfall in the region, topsoil moisture conditions are mostly at adequate levels.

Cropland topsoil moisture is:

- 2 per cent surplus;
- 66 per cent adequate;
- 27 per cent short; and
- 5 per cent very short.

Hayland topsoil moisture is:

- 2 per cent surplus;
- 68 per cent adequate;
- 29 per cent short; and
- 1 per cent very short.

Pasture topsoil moisture is:

- 2 per cent surplus;
- 65 per cent adequate;
- 31 per cent short; and
- 2 per cent very short.

While crops continue to mature in most places in the region, the recent cool weather conditions have slowed crop progress in some areas. Harvest is currently 1 per cent complete in the region. This is slightly below both the five-year and the 10-year average of 2 per cent but is on par with last year, when 1 per cent of the crop had been harvested at this time. Producers are looking forward to some much-needed moisture to help with grain filling in later seeded crops.

Winter cereals harvested:

- Fall rye: 92 per cent.

Spring cereals harvested:

- Triticale: 5 per cent; and
- Barley: 1 per cent.

Pulses harvested:

- Lentil: 2 per cent; and
- Field peas: 1 per cent.

Livestock producers are wrapping up haying and silaging. Average dryland alfalfa yields are 1.65 tonne (t) per acre, 1.35 t per acre for tame hay and 2.48 t per acre for greenfeed. Under irrigated systems, alfalfa yields average 2.02 t per acre while tame hay and greenfeed yields are 2.16 and 3.02 t per acre, respectively. Silage crops yielded an average of 5.30 t per acre.

Minor to moderate crop damage was reported in the region from hot, dry conditions. Minor damage from wind and from insects such as bertha armyworms were also reported. Some notable damage from aster yellows and alternaria black spot diseases were also reported.

West-Central Saskatchewan:

- Census Division 12 – Biggar, Delisle, Rosetown and Sonningdale areas
- Census Division 13 – Cut Knife, Kerrobert, Kindersley, Macklin, Plenty and Wilkie areas

This area experienced cool weather with light showers; however, conditions in the region remain dry. The cool conditions in some of the areas in the region slowed harvest and crop development. Harvest is slowly starting in the region with pulse crops being desiccated.

Rainfall in the region was variable with the Milton area receiving the most rain with 20 millimetres (mm), followed by the Mildred area with 16 mm and 15 mm in the Harris and Hillsdale areas.

The lack of significant rainfall impacted topsoil moisture levels. Topsoil moisture in the region is mostly rated as adequate with a notable proportion rated as short.

Cropland topsoil moisture is:

- 1 per cent surplus;
- 50 per cent adequate;
- 48 per cent short; and
- 1 per cent very short.

Hayland topsoil moisture is:

- 48 per cent adequate;
- 45 per cent short; and
- 7 per cent very short.

Pasture topsoil moisture is:

- 1 per cent surplus;
- 44 per cent adequate;
- 48 per cent short; and
- 7 per cent very short.

Although harvest has begun for a few crops, the amount harvested in the region currently represents less than 1 per cent of the total crop area. This is similar to last year when less than 1 per cent of crops were harvested at this time, but below the five-year average of 5 per cent and the 10-year average of 3 per cent.

Winter cereals harvested:

- Winter wheat: 10 per cent; and
- Fall rye: 7 per cent.

Spring cereals harvested:

- Barley: 8 per cent; and
- Oats: 3 per cent.

Oilseeds harvested:

- Mustard: 6 per cent.

Pulses harvested:

- Lentil: 1 per cent; and
- Field peas: 1 per cent.

With crops continuing to mature and pulse crop desiccation underway across the region, producers are looking forward to harvest operations getting into full swing in the coming weeks.

Haying and silaging in the region are largely wrapping up for the year. Dryland alfalfa yields averaged 1.82 tonne (t) per acre, while tame hay yielded 1.50 t per acre with greenfeed yield averaging 3.58 t per acre on average. Meanwhile, alfalfa yields under irrigation averaged 3 t per acre. Silage crops yielded 7.14 t per acre.

Minor crop damage in the region was caused by hot, dry conditions. Minor to moderate damage from bertha armyworms and diamondbacks were also reported, with reports of producers spraying for bertha armyworms. Sclerotinia and fusarium head blight diseases have also been reported to cause minor to moderate crop damage in the region.

Northeastern Saskatchewan:

- Census Division 14 – Choiceland, Hudson Bay, Kelvington, Melfort, and Nipawin areas
- Census Division 15 – Cudworth, Humboldt, Kinistino, Prince Albert, Rosthern and St. Brieux areas

Similar to most areas across the province, conditions in the region were mostly dry with cool temperatures welcomed in some areas as they helped reduce heat stress on some crops. With crops maturing and nearing harvest in some places, farmers are getting harvest machinery ready.

A few areas in the region received scattered showers, with the Arborfield and Nipawin areas receiving the highest precipitation amounts of 15 millimetres (mm). This was followed by the Barrier Valley and Duck Lake areas with 10 mm.

Topsoil moisture levels in the region remain adequate with a few rated as short.

Cropland topsoil moisture is:

- 1 per cent surplus;
- 72 per cent adequate; and
- 27 per cent short.

Hayland topsoil moisture is:

- 1 per cent surplus;
- 70 per cent adequate;
- 28 per cent short; and
- 1 per cent very short.

Pasture topsoil moisture is:

- 1 per cent surplus;
- 71 per cent adequate;
- 27 per cent short; and
- 1 per cent very short.

Harvest has yet to begin for most crops in the region. So far, only 1 per cent of barley and 34 per cent of fall rye have been harvested in the region, representing less than 1 per cent of total crops harvested. This is not far from both the five-year and the 10-year average of 1 per cent and not different from last year, when less than 1 per cent of the crop had been harvested at this time. With crops continuing to mature across the region, producers are looking forward to harvest operations getting into full swing in the coming weeks.

Dryland alfalfa yielded 2.08 tonne (t) per acre, with tame hay and greenfeed yield averaging 2.33 and 3.24 t per acre, respectively. Silage crops on the other hand yielded 6.50 t per acre.

Similar to most parts of the province, hot, dry weather conditions were reported as the major cause of crop damage in the northeast region. Minor to moderate crop damage from bertha armyworms were also reported with reports of producers spraying to manage these insects in some places in the region.

Northwestern Saskatchewan:

- Census Division 16 – Blaine Lake, Canwood, North Battleford, Radisson and Spiritwood areas
- Census Division 17 – Glaslyn, Maidstone, Meadow Lake, Pierceland and St. Walburg areas

Humid conditions in the region have caused crop diseases to appear in some areas. With crops continuing to advance in maturity, producers are looking forward to starting harvest soon. Although scattered showers in the region slowed haying operations, it was welcomed for some areas with later seeded crops.

The amount of rainfall varied with the Round Hill area receiving the most with 15 millimetres (mm), followed by the Parkdale and Loon Lake areas with 12 mm. The Meadow Lake area also received 10 mm of rain.

Topsoil moisture in the region is mostly rated as adequate.

Cropland topsoil moisture is:

- 1 per cent surplus;
- 92 per cent adequate;
- 6 per cent short; and
- 1 per cent very short.

Hayland topsoil moisture is:

- 1 per cent surplus;
- 93 per cent adequate;
- 5 per cent short; and
- 1 per cent very short.

Pasture topsoil moisture is:

- 1 per cent surplus;
- 94 per cent adequate; and
- 5 per cent short.


Most crops are still developing and are not ready to harvest. Currently, only 1 per cent each of lentils and field peas have been harvested, representing less than 1 per cent of total crop harvest in the region. This is similar to the five-year average and a year ago, when less than 1 per cent of crops in the region were harvested at this time. It is also not far off the 10-year average of 1 per cent.

With livestock producers wrapping up haying and silaging, dryland alfalfa yields have averaged 2.12 tonne (t) per acre in the region. Tame hay yielded 1.90 t per acre and greenfeed yielded 2.78 t per acre on average. Yields for silage crops averaged 6.67 t per acre.

Crop diseases such as sclerotinia, mycosphaerella blight, blackleg and ascochyta blight were reported to cause minor to severe crop damage in the region. Wind and aphids were also reported to cause minor crop damage in some areas.

Weekly Rainfall

from August 4 to August 10, 2026



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Weekly Rainfall Summary

(reported in millimeters)

1 inch=25 mm

for the period from August 4 to August 10, 2026

Census Division	RM No.	RM Name	Past Week	Since 1-Apr	Census Division	RM No.	RM Name	Past Week	Since 1-Apr	Census Division	RM No.	RM Name	Past Week	Since 1-Apr
1	3	ENNISKILLEN	0	287	8	138	WEBB	3	229	14	366	KELVINGTON	8	249
1	4	COALFIELDS	0	378	8	139	GULL LAKE	2	210	14	367	PONASS LAKE	N/A	306
1	32	RECIPROCITY	N/A	278	8	168	RIVERSIDE	0	193	14	394	HUDSON BAY	4	226
1	34	BROWNING	N/A	290	8	228	LACADENA	1	234	14	395	PORCUPINE	N/A	273
1	94	HAZELWOOD	N/A	0	8	229	MIRY CREEK	1	153	14	397	BARRIER VALLEY	10	194
1	95	GOLDEN WEST	N/A	225	8	231	HAPPYLAND	6	167	14	428	STAR CITY	4	226
1	2A	MOUNT PLEASANT	3	203	8	259	SNIE LAKE	5	194	14	456	ARBORFIELD	15	272
2	10	HAPPY VALLEY	0	196	8	138A	WEBB	3	156	14	457	CONNAUGHT	N/A	67
2	38	LAURIER	0	248	8	228A	LACADENA	N/A	64	14	486	MOOSE RANGE	N/A	212
2	66	GRIFFIN	0	294	8	257A	MONET	0	188	14	487	NIPAWIN	15	242
2	67	WEYBURN	N/A	101	8	259A	SNIE LAKE	N/A	141	14	488	TORCH RIVER	2	227
2	70	KEY WEST	N/A	12	9	241	CALDER	2	385	14	394A	HUDSON BAY	N/A	245
2	100	ELMSTHORPE	0	209	9	243	WALLACE	0	232	14	397A	BARRIER VALLEY	9	244
2	100A	ELMSTHORPE	0	223	9	245	GARRY	1	159	14	488A	TORCH RIVER	0	116
2	38A	LAURIER	0	215	9	273	SLIDING HILLS	0	318	14	488B	TORCH RIVER	N/A	5
3	73	STONEHENGE	0	209	9	274	GOOD LAKE	0	244	15	369	ST. PETER	N/A	208
3	74	WOOD RIVER	0	207	9	301	ST. PHILIPS	N/A	143	15	371	BAYNE	6	252
3	75	PINTO CREEK	0	145	9	305	INVERMAY	0	310	15	372	GRANT	0	182
3	76	AUVERGNE	0	199	9	331	LIVINGSTON	0	238	15	373	ABERDEEN	N/A	186
3	101	TERRELL	N/A	0	9	241A	CALDER	4	474	15	399	LAKE LENORE	3	238
3	102	LAKE JOHNSTON	0	195	9	245A	GARRY	2	213	15	400	THREE LAKES	N/A	311
3	106	WHISKA CREEK	0	178	10	246	ITUNA BON ACCORD	0	131	15	402	FISH CREEK	5	176
3	74A	KEY WEST	0	170	10	247	KELLROSS	N/A	216	15	429	FLETT'S SPRINGS	0	234
4	51	RENO	0	130	10	248	TOUCHWOOD	1	204	15	430	INVERGORDON	4	124
4	79	ARLINGTON	0	218	10	277	EMERALD	2	208	15	459	KINISTINO	3	262
4	110	PIAPOT	N/A	217	10	279	MOUNT HOPE	2	191	15	461	PRINCE ALBERT	N/A	224
4	77A	WISE CREEK	0	152	10	307	ELFROS	0	399	15	463	DUCK LAKE	10	246
4	78A	GRASSY CREEK	0	127	10	309	PRAIRIE ROSE	0	255	15	491	BUCKLAND	3	108
4	79A	ARLINGTON	0	236	10	336	SASMAN	0	299	15	520	PADDOCKWOOD	N/A	20
5	122	MARTIN	5	262	10	339	LEROY	1	237	15	521	LAKELAND	N/A	20
5	124	KINGSLEY	6	257	10	246A	ITUNA BON ACCORD	N/A	105	15	371A	BAYNE	0	176
5	151	ROCANVILLE	2	241	10	248A	TOUCHWOOD	2	347	15	371B	BAYNE	0	233
5	155	WOLSELEY	6	202	10	276A	FOAM LAKE	N/A	341	15	399A	LAKE LENORE	N/A	16
5	181	LANGENBURG	1	424	10	276B	FOAM LAKE	2	456	15	403A	ROSTHERN	0	249
5	183	FERTILE BELT	0	278	10	276C	FOAM LAKE	3	358	15	403C	ROSTHERN	2	247
5	211	CHURCHBRIDGE	0	296	10	276D	FOAM LAKE	0	260	15	461A	PRINCE ALBERT	0	192
5	213	SALT COATS	0	329	10	277A	EMERALD	N/A	270	15	461B	PRINCE ALBERT	N/A	2
5	215	STANLEY	0	218	10	279A	MOUNT HOPE	N/A	158	16	434	BLAINE LAKE	4	151
5	125A	CHESTERFIELD	20	208	11	251	BIG ARM	N/A	232	16	435	REDBERRY	N/A	295
5	154A	ELCAPO	N/A	213	11	280	WREFORD	N/A	83	16	436	DOUGLAS	0	192
5	183A	FERTILE BELT	0	332	11	283	ROSEDALE	6	260	16	437	NORTH BATTLEFORD	5	342
5	211A	CHURCHBRIDGE	0	317	11	284	RUDY	9	74	16	466	MEETING LAKE	7	301
6	127	FRANCIS	3	221	11	314	DUNDURN	2	203	16	467	ROUND HILL	5	259
6	128	LAJORD	0	111	11	344	CORMAN PARK	N/A	154	16	493	SHELLBROOK	N/A	174
6	130	REDBURN	0	238	11	282A	McCRANEY	N/A	155	16	494	CANWOOD	0	116
6	156	INDIAN HEAD	3	246	12	286	MILDEN	16	221	16	497	MEDSTEAD	N/A	267
6	160	PENSE	N/A	160	12	287	ST. ANDREWS	N/A	160	16	437A	NORTH BATTLEFORD	N/A	279
6	186	ABERNETHY	N/A	231	12	288	PLEASANT VALLEY	8	163	16	467A	ROUND HILL	15	322
6	190	DUFFERIN	0	178	12	316	HARRIS	15	148	17	468	MEOTA	0	215
6	216	TULLYMET	N/A	10	12	317	MARRIOTT	5	208	17	470	PAYNTON	N/A	77
6	217	LIPTON	2	331	12	345	VANSCOY	5	135	17	471	ELDON	2	276
6	219	LONGLAKETON	4	242	12	346	PERDUE	8	183	17	498	PARKDALE	12	322
6	221	SARNIA	4	217	12	347	BIGGAR	0	239	17	499	MERVIN	2	232
6	127B	FRANCIS	N/A	116	12	376	EAGLE CREEK	0	225	17	502	BRITANNIA	1	192
6	159A	SHERWOOD	5	136	12	377	GLENSIDE	1	208	17	561	LOON LAKE	12	175
6	159B	SHERWOOD	2	191	12	285A	FERTILE VALLEY	5	149	17	588	MEADOW LAKE	10	328
6	160A	PENSE	3	142	13	290	KINDERSLEY	N/A	166	17	622	BEAVER RIVER	9	274
6	190A	DUFFERIN	5	125	13	292	MILTON	20	353	17	498A	PARKDALE	7	269
6	190B	DUFFERIN	3	207	13	321	PRAIRIEDALE	10	255	17	501A	FRENCHMAN BUTTE	4	243
6	190C	DUFFERIN	5	251	13	350	MARIPOSA	8	208	17	501C	FRENCHMAN BUTTE	N/A	50
6	216A	TULLYMET	N/A	75	13	351	PROGRESS	9	238	17	561A	LOON LAKE	4	178
6	217A	LIPTON	N/A	2	13	379	REFORD	N/A	93					
6	219A	LONGLAKETON	0	226	13	382	EYE HILL	0	217					
6	219B	LONGLAKETON	N/A	60	13	409	BUFFALO	0	270					
6	220A	McKILLOP	3	164	13	440	HILLSDALE	15	256					
6	220B	McKILLOP	1	1	13	442	MANITOU LAKE	0	232					
7	132	HILLSBOROUGH	1	300	13	292A	MILTON	6	177					
7	136	COULEE	N/A	151	13	320A	OAKDALE	11	178					
7	161	MOOSE JAW	1	143	13	321A	PRAIRIEDALE	N/A	35					
7	162	CARON	2	211	13	409A	BUFFALO	1	228					
7	165	MORSE	0	250										
7	193	EYEBROW	3	245										
7	223	HURON	1	186										
7	132A	HILLSBOROUGH	0	118										
7	136A	COULEE	N/A	0										
7	161A	MOOSE JAW	N/A	38										
7	162A	CARON	5	188										
7	222A	CRAIK	3	230										
7	223A	HURON	4	181										


Municipality No: A, B, C and D - more than one reporter

These precipitation amounts represent point locations within each municipality and do not necessarily reflect the whole R. M.

N/A indicates that rainfall was not reported for the week


Cumulative Rainfall

from April 1 to August 10, 2026



Cropland Topsoil Moisture Conditions

from August 4 to August 10, 2026



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Topsoil Moisture Conditions-for the period of August 4th to August 10th, 2026


Cropland				
Region	Surplus	Adequate	Short	Very Short
South East	0%	43%	55%	2%
South West	0%	18%	58%	24%
East Central	2%	66%	27%	5%
West Central	1%	50%	48%	1%
North East	1%	72%	27%	0%
North West	1%	92%	6%	1%
Provincial Average	1%	49%	42%	8%

Hayland				
Region	Surplus	Adequate	Short	Very Short
South East	0%	42%	55%	3%
South West	0%	19%	54%	27%
East Central	2%	68%	29%	1%
West Central	0%	48%	45%	7%
North East	1%	70%	28%	1%
North West	1%	93%	5%	1%
Provincial Average	1%	49%	42%	8%

Pasture				
Region	Surplus	Adequate	Short	Very Short
South East	0%	36%	59%	5%
South West	0%	13%	60%	27%
East Central	2%	65%	31%	2%
West Central	1%	44%	48%	7%
North East	1%	71%	27%	1%
North West	1%	94%	5%	0%
Provincial Average	1%	39%	46%	14%

Hay Topsoil Moisture Conditions


from August 4 to August 10, 2026



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Pasture Topsoil Moisture Conditions


from August 4 to August 10, 2026




NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Maximum Temperature

from August 4 to August 10, 2026



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.




Projection: UTM Zone 13 Datum: NAD83



Data Sources:	
Temperature data - Saskatchewan Ministry of Environment (Wildfire Management Branch) and Environment Canada.	
Temperature data compiled and quality controlled by Agriculture and Agri-Food Canada	
IDW interpolation (power 3.5, fixed radius 300 km)	
Geomatics Services, Ministry of Agriculture	August 12, 2022

Minimum Temperature

from August 4 to August 10, 2026



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.