

Diabetes Management in Chronic Kidney Disease: A Consensus Report by the American Diabetes Association (ADA) and Kidney Disease: Improving Global Outcomes (KDIGO)

Diabetes Care 2022;45:3075–3090.

				Albuminuria categories		
				Description and range		
				A1	A2	A3
CKD is classified based on:				Normal to mildly increased	Moderately increased	Severely increased
• Cause (C)						
• GFR (G)						
• Albuminuria (A)				<30 mg/g <3 mg/mmol	30–299 mg/g 3–29 mg/mmol	≥300 mg/g ≥30 mg/mmol
GFR categories (mL/min/1.73 m ²)	Description and range					
	G1	Normal or high	≥90	Screen 1	Treat 1	Treat and refer 3
	G2	Mildly decreased	60–89	Screen 1	Treat 1	Treat and refer 3
	G3a	Mildly to moderately decreased	45–59	Treat 1	Treat 2	Treat and refer 3
	G3b	Moderately to severely decreased	30–44	Treat 2	Treat and refer 3	Treat and refer 3
	G4	Severely decreased	15–29	Treat and refer* 3	Treat and refer* 3	Treat and refer 4+
	G5	Kidney failure	<15	Treat and refer 4+	Treat and refer 4+	Treat and refer 4+

Low risk (if no other markers of kidney disease, no CKD)

Moderately increased risk

High risk

Very high risk

Figure 2—Risk of CKD progression, frequency of visits, and referral to nephrology according to GFR and albuminuria. The numbers in the boxes are a guide to the frequency of screening or monitoring (number of times per year). Green reflects no evidence of CKD by eGFR or albuminuria, with screening indicated once per year. For monitoring of prevalent CKD, suggested monitoring varies from once per year (yellow) to four times or more per year (i.e., every 1–3 months, [deep red]) according to risks of CKD progression and CKD complications. These are general parameters only, based on expert opinion, and underlying comorbid conditions and disease state must be taken into account, as well as the likelihood of impacting a change in management for any individual patient. CKD, chronic kidney disease; GFR, glomerular filtration rate.