Crop Report

For the Period May 28 to June 3, 2019

Published by the Ministry of Agriculture ISSN 0701 7085 Report number 06, June 6, 2019

Seeding operations are mostly complete across the province, but there are a few fields still being seeded for green feed and silage. All six crop districts in the province are reporting that 99 per cent of the crop is in the ground, compared to 92 per cent last week and the five-year average (2014-2018) of 93 per cent for this time of year. Ninety-nine per cent of the spring wheat, durum, canola, canary seed are seeded, while 98 per cent of the oats, barley and flax are planted.

Little rainfall was reported this past week throughout the province, with amounts ranging from nothing to 18 mm in the Rocanville area. Significant rain is needed soon across the province to help with crop growth and emergence as well as with the establishment and growth of hay land and pastures.

Per cent seeded all crops June 3, 2019									
Spring wheat	99								
Durum	99								
Barley	98								
Oats	98								
Canary seed	99								
Flax	98								
Canola	99								
Mustard	92								
Soybeans	100								
Lentils	100								
Field peas	100								
Chickpeas	100								

Topsoil moisture conditions continue to deteriorate across the province due to strong winds and the lack of moisture. Cropland topsoil moisture is rated as 22 per cent adequate, 47 per cent short and 31 per cent very short. Topsoil moisture on hay land and pasture is rated as 18 per cent adequate, 42 per cent short and 40 per cent very short. Hay and pasture growth has been slow due to little rainfall.

Seeding Progress in SK Per cent seeded **All Crops** June 3, 2019 99 June 4, 2018 96 June 5, 2017 94 June 6, 2016 98 97 June 1, 2015 June 2, 2014 78 5 year avg. 93 (2014-2018)10 year avg. 88

(2009-2018)

One year ago

Ninety-six per cent of the crop

had been seeded. The majority of the province received much-

needed moisture, which

helped to alleviate concerns

about dry field conditions.

Follow the 2019 Crop Report

The majority of crop conditions range from poor to good across the province. Crop growth is delayed in much of the province due to the lack of moisture. The majority of crop damage this past week was due to strong winds, insects such as flea beetles and cutworms, and the lack of moisture. Farmers are busy making in-crop pesticide applications to control weeds and insects.

For further information, contact Cory Jacob, PAg, Crops Extension Specialist, Regional Services Branch, Toll Free: 1-866-457-2377 or 306-694-3587, Email: cropreport@gov.sk.ca. Also available on the Ministry of Agriculture website at saskatchewan.ca/crop-report.







Southeastern Saskatchewan:

- Crop District 1 Carnduff, Estevan, Redvers, Moosomin and Kipling areas
- Crop District 2 Weyburn, Milestone, Moose Jaw, Regina and Qu'Appelle areas
- Crop District 3ASE Radville, Minton and Lake Alma areas

Seeding is essentially complete in the region as 99 per cent of the crop is now in the ground. This is up from ninety-four per cent last week and ahead of the five-year (2014-2018) average of 92 per cent for this time of year. Ninety-six per cent of the oats and 99 per cent of the flax, canola, barley and wheat are planted.

Crop District	% seeded (June 3, 2019)
1A	99
1B	99
2A	99
2B	99
3ASE	99
Region average	99

Small amounts of rainfall were reported in the region this past week, with 3 mm falling in the Glenavon and Broadview areas and 7 mm in the Tantallon area, while 6 mm was reported in the Grenfell area. The Indian Head area has received the most precipitation since April 1 (64 mm). Significant rainfall is needed in much of the region to help crop germination as well as hay and pasture growth.

Topsoil moisture conditions have continued to deteriorate due to strong winds, warm temperatures and a lack of rainfall. Many areas remain in need of significant moisture. Cropland topsoil moisture is rated as 24 per cent adequate, 51 per cent short and 25 per cent very short. Hay land and pasture topsoil moisture is rated as 13 per cent adequate, 49 per cent short and 38 per cent very short. Rain will be needed soon to help crops germinate and emerge and pastures grow. Crop District 2A is reporting that 50 per cent of cropland and 58 per cent of hay land and pasture are very short topsoil moisture at this time. Yields will likely be compromised without substantial rain in the coming weeks.

Crops are slow to emerge, patchy and behind their normal developmental stages for this time of year. The majority of crops across the region are in poor to good condition. There have been reports of thin and stunted winter cereal crops that are prematurely heading-out due to the lack of moisture. Other crops are at a stand-still in the field and many producers may delay in-crop herbicide applications until rain comes. Flea beetles and cutworms are damaging canola crops, while strong winds and a lack of rain also caused crop damage this past week.

Farmers are busy finishing seeding, controlling weeds and picking rocks.

Southwestern Saskatchewan:

- Crop District 3ASW Coronach, Assiniboia and Ogema areas
- Crop District 3AN Gravelbourg, Mossbank, Mortlach and Central Butte areas
- Crop District 3B Kyle, Swift Current, Shaunavon and Ponteix areas
- Crop District 4 Consul, Maple Creek and Leader areas

The southwestern region has essentially completed seeding operations as 99 per cent of the crop is now seeded, up from 96 per cent last week and ahead of the five-year (2014-2018) average of 94 per cent for this time of year. Ninety-seven per cent of the oats, barley and canola are seeded, while 99 per cent of the durum and 98 per cent of the flax are in the ground.

Only the Consul area reported rainfall this past week, receiving 0.8 mm. The Shaunavon area has received the most precipitation since April 1 (92.5 mm).

Strong winds and warm weather continue to cause topsoil moisture conditions to deteriorate in the region. Significant rainfall is needed to help crop germination and growth as well as pasture and hay

Crop District	% Seeded (June 3, 2019)
3ASW	99
3AN	99
3BS	99
3BN	98
4A	99
4B	99
Region average	99

growth. Cropland topsoil moisture is rated as 20 per cent adequate, 53 per cent short and 27 per cent very short. Hay land and pasture topsoil moisture is rated as 15 per cent adequate, 43 per cent short and 42 per cent very short. Crop District 3BN is reporting that 50 per cent of the cropland and 70 per cent of the hay land and pasture are very short top soil moisture at this time. Significant yield loss will result if substantial rainfall is not received in the coming weeks.

Crop emergence has been slow and patchy, and crops are behind in their growth and development due to the lack of moisture. The lack of moisture is also causing some thin and stunted winter cereal crops to prematurely head-out. Overall, crop conditions are mostly poor to good. Most of the crop damage this past week was caused by strong winds, flea beetles and a lack of moisture.

Concerns remain in the area over poor quality livestock water. Farmers are busy finishing seeding and controlling weeds.

East-Central Saskatchewan:

- Crop District 5 Melville, Yorkton, Cupar, Kamsack, Foam Lake, Preeceville and Kelvington areas
- Crop District 6A Lumsden, Craik, Watrous and Clavet areas

Seeding has basically wrapped up in the region, but there may yet be fields seeded to greenfeed or silage. Ninety-nine per cent of the is now in the ground, up from 91 per cent last week and well ahead of the five-year (2014-2018) average of 90 per cent for this time of year. Ninety-eight per cent

Crop District	% Seeded (June 3, 2019)
5A	99
5B	99
6A	99
Region average	99

of the oats, 97 per cent of the flax, 99 per cent of the spring wheat and canola are seeded.

The region received little rainfall last week, with most areas reporting no rain. Eighteen mm fell in the Rocanville area; the Lipton and Rhein areas reported 1 mm, while 3 mm fell in the Rama area and 9.7 mm in the Ituna area. The Rocanville area has received the most precipitation since April 1 (45 mm).

Topsoil moisture conditions continue to worsen due to the strong winds and lack of moisture. Substantial rainfall is needed across the region to help crop germination and growth as well as pasture growth. Cropland topsoil moisture is rated as 14 per cent adequate, 39 per cent short and 47 per cent very short. Hay land and pasture topsoil moisture is rated as nine per cent adequate, 33 per cent short and 58 per cent very short. Crop District 6A is reporting that 71 per cent of cropland and 78 per cent of hay land and pasture are very short topsoil moisture at this time. Timely rains are needed in the coming weeks to prevent significant yield loss.

Crop emergence has been and patchy slow due to the lack of moisture; this has also put the majority of crops behind their normal developmental stages for this time of year. The majority of crops are in poor to good condition. Crop damage this past week was caused by frost, dry conditions, strong winds, flea beetles and cutworms.

Farmers are busy completing seeding, controlling weeds and moving cattle to pasture.

West-Central Saskatchewan:

- Crop District 6B Hanley, Outlook, Loreburn, Saskatoon and Arelee areas
- Crop District 7A Rosetown, Kindersley, Eston and Major areas
- Crop District 7B Kerrobert, Macklin, Wilkie and Biggar areas

Ninety-nine per cent of the acres are now seeded in the region, up from 92 per cent last week. This is ahead of the five-year (2014-2018) average of 94 per cent for this time of year. Ninety-two per cent of the oats, 96 per cent of the barley, 98 per cent of the flax and spring wheat, and 99 per cent of the

Crop District	% Seeded (June 3, 2019)
6B	97
7A	99
7B	99
Region average	99

canola and soybeans are now seeded. There have been reports that some producers are discontinuing seeding operations due to a lack of moisture and will resume them if rain falls in the coming weeks.

Only the Semans area reported rainfall this past week, receiving 0.1 mm. The Landis area has received the most precipitation since April 1 (43 mm).

Topsoil moisture conditions have continued to deteriorate in the region with the strong winds and lack of rainfall. Cropland topsoil moisture is rated as seven per cent adequate, 51 per cent short and 42 per cent very short. Hay land and pasture topsoil moisture is rated as four per cent adequate, 34 per cent short and 62 per cent very short. Crop District 7A is reporting that 52 per cent of cropland and 65 per cent of hay land and pasture are very short topsoil moisture at this time, while Crop District 7B is reporting that 44 per cent of cropland and 70 per cent of hay land and pasture are very short topsoil moisture. Significant yield loss will result if substantial rainfall is not received in the coming weeks.

The dry conditions have resulted in slow crop emergence and growth as well as slow hay and pasture growth. The majority of crops are in poor to good condition. Crop damage this past week was from dry conditions, wind, flea beetles and cutworms. There have been reports of grasshoppers damaging forage crops and pastures in the region as well.

Producers remain concerned about the dry soil moisture conditions and livestock water quality. Farmers are busy finishing seeding and moving cattle.

Northeastern Saskatchewan:

- Crop District 8 Hudson Bay, Tisdale, Melfort, Carrot River, Humboldt, Kinistino, Cudworth and Aberdeen areas
- Crop District 9AE Prince Albert, Choiceland and Paddockwood areas

The northeastern region has basically completed seeding operations with 99 per cent of the crop now seeded, which is up from 90 per cent last week and ahead of the five-year (2014-2018) average of 91 per cent for this time of year. Ninety-nine per cent of the barley, oats, flax, canola and spring wheat are in the ground.

Crop District	% Seeded (June 3, 2019)
8A	99
8B	99
9AE	99
Region average	99

Varying amounts of rainfall were reported last week, ranging from nothing to 16 mm in the Garrick area. The Hudson Bay area reported 10 mm of rain, the Bjorkdale area 8 mm, the Star City area 6 mm and the Arborfield and Humboldt areas 4 mm. The Garrick area has received the most precipitation since April 1 (58 mm).

Topsoil moisture conditions have continued to deteriorate due to strong winds and warm temperatures. Cropland topsoil moisture is rated as 43 per cent adequate, 44 per cent short and 13 per cent very short. Hay land and pasture topsoil moisture is rated as 46 per cent adequate, 43 per cent short and 11 per cent very short. Crop reporters have indicated that rain is welcomed at any time to encourage crop germination and growth as well as hay and pasture growth.

Overall, crops are at or behind their normal developmental stages for this time of the year. Emergence has been delayed in many areas by a lack of moisture. Crops are mostly in poor to good condition. The majority of crop damage this past week was caused by strong winds, flea beetles, cutworms and the lack of moisture.

Farmers are busy completing seeding operations, controlling weeds and moving cattle to pasture.

Northwestern Saskatchewan:

- Crop District 9AW Shellbrook, North Battleford, Big River and Hafford areas
- Crop District 9B Meadow Lake, Turtleford, Pierceland, Maidstone and Lloydminster areas

Producers in the region have the 2019 crop mostly seeded. Ninety-nine per cent of the crop is in the ground, up significantly from 87 per cent last week and ahead of the five-year (2014-2018) average of 93 per cent for this time of year. Ninety-seven per

Crop District	% Seeded (June 3, 2019)
9AW	97
9B	99
Region average	99

cent of the oats and barley, 93 per cent of the flax and 99 per cent of the spring wheat and canola are seeded.

The region received little rainfall last week, with most areas reporting no rain to 5 mm in the Meadow Lake area. The Radisson, Hafford, Mayfair, Lloydminster and Barthel areas reported 2 mm. The Turtleford area has received the most precipitation since April 1 (101.5 mm).

As in most of the province, topsoil moisture conditions have worsened and rainfall would be welcomed across the region. Cropland topsoil moisture is rated as 47 per cent adequate, 39 per cent short and 14 per cent very short. Hay land and pasture topsoil moisture is rated as 36 per cent adequate, 48 per cent short and 16 per cent very short. Significant rainfall is needed throughout the region to even out crop germination and growth and would be highly beneficial to hay crops and pasture.

Generally, crop growth is slow and emergence patchy, especially on the shallow-seeded crops. Lack of moisture has slowed cropland and hay land and pasture progress. Most crops in the are in poor to good condition. Most of the crop damage this past week was caused by strong winds, flea beetles, cutworms and the lack of moisture.


Farmers are busy finishing seeding, controlling weeds and moving cattle to pasture.

Crop Conditions - June 3, 2019


Winter wheat Fall rye Spring wheat Durum Oat Barley Canaryseed											
% excellent 3 1 3 2 3 2 0 % good 31 25 33 31 43 31 11 % fair 37 57 45 55 34 40 51 % poor 28 14 15 11 18 24 36 % very poor 1 3 4 1 2 3 2 Flax Canola Mustard Soybean Pea Lentil Chickpea % excellent 0 2 0 0 3 1 0 % good 19 21 0 37 46 35 33 % fair 49 43 40 52 41 52 67 % poor 28 26 59 11 9 11 0 % excellent 5 0 2 1 3 1 0 % excellent											
% good 31 25 33 31 43 31 11 % fair 37 57 45 55 34 40 51 % poor 28 14 15 11 18 24 36 % very poor 1 3 4 1 2 3 2 Flax Canola Mustard Soybean Pea Lentil Chickpea % excellent 0 2 0 0 3 1 0 % good 19 21 0 37 46 35 33 % fair 49 43 40 52 41 52 67 % poor 28 26 59 11 9 11 0 % very poor 4 8 1 0 1 1 0 * excellent 5 0 2 1 3 1 0 % good											
% fair 37 57 45 55 34 40 51 % poor 28 14 15 11 18 24 36 % very poor 1 3 4 1 2 3 2 Flax Canola Mustard Soybean Pea Lentil Chickpea % excellent 0 2 0 0 3 1 0 % good 19 21 0 37 46 35 33 % fair 49 43 40 52 41 52 67 % poor 28 26 59 11 9 11 0 % very poor 4 8 1 0 1 1 0 Southeast Winter wheat Fall rye Spring wheat Durum Oat Barley Canaryseed % excellent 5 0 2 1 3											
% poor 28 14 15 11 18 24 36 % very poor 1 3 4 1 2 3 2 Flax Canola Mustard Soybean Pea Lentil Chickpea % excellent 0 2 0 0 3 1 0 % good 19 21 0 37 46 35 33 % fair 49 43 40 52 41 52 67 % poor 28 26 59 11 9 11 0 % very poor 4 8 1 0 1 1 0 % excellent 5 0 2 1 3 1 0 % excellent 5 0 2 1 3 1 0 % good 27 36 50 24 67 33 11 % fair 38 54 41 67 25 46 66 % poor <											
% very poor 1 3 4 1 2 3 2 Flax Canola Mustard Soybean Pea Lentil Chickpea % excellent 0 2 0 0 3 1 0 % good 19 21 0 37 46 35 33 % fair 49 43 40 52 41 52 67 % poor 28 26 59 11 9 11 0 % very poor 4 8 1 0 1 1 0 Southeast Winter wheat Fall rye Spring wheat Durum Oat Barley Canaryseed % excellent 5 0 2 1 3 1 0 % good 27 36 50 24 67 33 11 % fair 38 54 41 67 25											
Flax Canola Mustard Soybean Pea Lentil Chickpea % excellent 0 2 0 0 3 1 0 % good 19 21 0 37 46 35 33 % fair 49 43 40 52 41 52 67 % poor 28 26 59 11 9 11 0 % very poor 4 8 1 0 1 1 0 Southeast Winter wheat Fall rye Spring wheat Durum Oat Barley Canaryseed % excellent 5 0 2 1 3 1 0 % good 27 36 50 24 67 33 11 % fair 38 54 41 67 25 46 66 % poor 28 9 6 7 5											
% excellent 0 2 0 0 3 1 0 % good 19 21 0 37 46 35 33 % fair 49 43 40 52 41 52 67 % poor 28 26 59 11 9 11 0 % very poor 4 8 1 0 1 1 0 Southeast Winter wheat Fall rye Spring wheat Durum Oat Barley Canaryseed % excellent 5 0 2 1 3 1 0 % good 27 36 50 24 67 33 11 % fair 38 54 41 67 25 46 66 % poor 28 9 6 7 5 17 21 % very poor 2 1 1 1 0											
% good 19 21 0 37 46 35 33 % fair 49 43 40 52 41 52 67 % poor 28 26 59 11 9 11 0 Southeast Winter wheat Fall rye Spring wheat Durum Oat Barley Canaryseed % excellent 5 0 2 1 3 1 0 % good 27 36 50 24 67 33 11 % fair 38 54 41 67 25 46 66 % poor 28 9 6 7 5 17 21 % very poor 2 1 1 1 0 3 2 Flax Canola Mustard Soybean Pea Lentil Chickpea % excellent 0 0 1 0 0											
% fair 49 43 40 52 41 52 67 % poor 28 26 59 11 9 11 0 % very poor 4 8 1 0 1 1 0 Southeast Winter wheat Fall rye Spring wheat Durum Oat Barley Canaryseed % excellent 5 0 2 1 3 1 0 % good 27 36 50 24 67 33 11 % fair 38 54 41 67 25 46 66 % poor 28 9 6 7 5 17 21 % very poor 2 1 1 1 0 3 2 Flax Canola Mustard Soybean Pea Lentil Chickpea % excellent 0 0 0 0 0 % good 23 27 50 42 47 39 30<											
% poor 28 26 59 11 9 11 0 % very poor 4 8 1 0 1 1 0 Southeast Winter wheat Fall rye Spring wheat Durum Oat Barley Canaryseed % excellent 5 0 2 1 3 1 0 % good 27 36 50 24 67 33 11 % fair 38 54 41 67 25 46 66 % poor 28 9 6 7 5 17 21 % very poor 2 1 1 1 0 3 2 Flax Canola Mustard Soybean Pea Lentil Chickpea % excellent 0 0 0 0 0 0 % good 23 27 50 42 47 39											
% very poor 4 8 1 0 1 1 0 Southeast Winter wheat Fall rye Spring wheat Durum Oat Barley Canaryseed % excellent 5 0 2 1 3 1 0 % good 27 36 50 24 67 33 11 % fair 38 54 41 67 25 46 66 % poor 28 9 6 7 5 17 21 % very poor 2 1 1 1 0 3 2 Flax Canola Mustard Soybean Pea Lentil Chickpea % excellent 0 0 0 0 0 % good 23 27 50 42 47 39 30 % fair 62 45 44 57 43 49 70 </td											
Southeast Winter wheat Fall rye Spring wheat Durum Oat Barley Canaryseed % excellent 5 0 2 1 3 1 0 % good 27 36 50 24 67 33 11 % fair 38 54 41 67 25 46 66 % poor 28 9 6 7 5 17 21 % very poor 2 1 1 1 0 3 2 Flax Canola Mustard Soybean Pea Lentil Chickpea % excellent 0 0 1 0 0 0 0 % good 23 27 50 42 47 39 30 % fair 62 45 44 57 43 49 70											
Winter wheat Fall rye Spring wheat Durum Oat Barley Canaryseed % excellent 5 0 2 1 3 1 0 % good 27 36 50 24 67 33 11 % fair 38 54 41 67 25 46 66 % poor 28 9 6 7 5 17 21 % very poor 2 1 1 1 0 3 2 Flax Canola Mustard Soybean Pea Lentil Chickpea % excellent 0 0 1 0 0 0 0 % good 23 27 50 42 47 39 30 % fair 62 45 44 57 43 49 70											
% excellent 5 0 2 1 3 1 0 % good 27 36 50 24 67 33 11 % fair 38 54 41 67 25 46 66 % poor 28 9 6 7 5 17 21 % very poor 2 1 1 1 0 3 2 Flax Canola Mustard Soybean Pea Lentil Chickpea % excellent 0 0 1 0 0 0 0 % good 23 27 50 42 47 39 30 % fair 62 45 44 57 43 49 70											
% good 27 36 50 24 67 33 11 % fair 38 54 41 67 25 46 66 % poor 28 9 6 7 5 17 21 % very poor 2 1 1 1 0 3 2 Flax Canola Mustard Soybean Pea Lentil Chickpea % excellent 0 0 1 0 0 0 0 % good 23 27 50 42 47 39 30 % fair 62 45 44 57 43 49 70											
% fair 38 54 41 67 25 46 66 % poor 28 9 6 7 5 17 21 % very poor 2 1 1 1 0 3 2 Flax Canola Mustard Soybean Pea Lentil Chickpea % excellent 0 0 1 0 0 0 0 % good 23 27 50 42 47 39 30 % fair 62 45 44 57 43 49 70											
% poor 28 9 6 7 5 17 21 % very poor 2 1 1 1 0 3 2 Flax Canola Mustard Soybean Pea Lentil Chickpea % excellent 0 0 1 0 0 0 0 % good 23 27 50 42 47 39 30 % fair 62 45 44 57 43 49 70											
% very poor 2 1 1 1 0 3 2 Flax Canola Mustard Soybean Pea Lentil Chickpea % excellent 0 0 1 0 0 0 0 % good 23 27 50 42 47 39 30 % fair 62 45 44 57 43 49 70											
Flax Canola Mustard Soybean Pea Lentil Chickpea % excellent 0 0 1 0 0 0 0 % good 23 27 50 42 47 39 30 % fair 62 45 44 57 43 49 70											
% excellent 0 0 1 0 0 0 0 % good 23 27 50 42 47 39 30 % fair 62 45 44 57 43 49 70											
% good 23 27 50 42 47 39 30 % fair 62 45 44 57 43 49 70											
% fair 62 45 44 57 43 49 70											
% very poor 2 11 1 0 1 1 0											
Southwest											
Winter wheat Fall rye Spring wheat Durum Oat Barley Canaryseed											
% excellent 4 5 2 1 2 8 0											
% good 14 16 26 36 42 23 19											
% fair 77 73 57 53 54 64 52											
% poor 5 6 15 8 2 5 29											
% very poor 0 0 0 1 0 0											
Flax Canola Mustard Soybean Pea Lentil Chickpea											
% excellent 0 0 0 0 1 2 0											
% good 45 20 0 0 55 38 34											
% fair 51 63 44 30 40 56 66											
% poor 4 16 56 70 4 4 0											
% very poor 0 1 0 0 0 0											
East-central											
Winter wheat Fall rye Spring wheat Durum Oat Barley Canaryseed											
% excellent 0 0 0 0 0 0											
% good 43 0 19 10 23 15 1											
% fair 25 63 48 48 37 44 97											
% poor 31 34 25 22 36 35 2											
% very poor 1 3 8 20 4 6 0											

East-central (continued)											
Flax Canola Mustard Soybean Pea Lentil Chickpe											
% excellent	0	0	0	0	0	0	N/A				
% good	5	7	31	19	28	35	N/A				
% fair	25	34	2	26	51	37	N/A				
% poor	61	41	67	55	21	28	N/A				
% very poor	9	18	0	0	0	0	N/A				
West-central											
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed				
% excellent	0	0	2	0	0	0	0				
% good	0	5	25	15	25	23	0				
% fair	48	45	46	32	56	34	42				
% poor	52	33	21	50	14	39	56				
% very poor	0	17	6	3	5	4	2				
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea				
% excellent	0	1	0	0	1	0	0				
% good	13	16	0	0	39	26	0				
% fair	39	38	32	60	41	50	0				
% poor	47	39	66	40	16	23	100				
% very poor	1	6	2	0	3	1	0				
Northeast											
	Winter wheat	Fall rye	Spring wheat		Oat	Barley	Canaryseed				
% excellent	0	0	5	2	6	5	2				
% good	62	40	54	42	61	53	52				
% fair	26	45	36	42	29	35	39				
% poor	12	15	4	12	3	6	5				
% very poor	0	0	1	2	1	1	2				
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea				
% excellent	1	4	0	0	9	0	N/A				
% good	41	40	15	8	57	15	N/A				
% fair	41	38	50	70	29	50	N/A				
% poor	16	17	25	18	4	25	N/A				
% very poor	1	1	10	4	1	10	N/A				
	140	- "	Northwe								
0/ 01/5-11/5-1	Winter wheat	Fall rye	Spring wheat		Oat	Barley	Canaryseed				
% excellent	0	0	13	N/A	6	5	N/A				
% good	100	37	40	N/A	46	47	N/A				
% fair	0	27	40	N/A	38	39	N/A				
% poor	0	3	7	N/A	10	9	N/A				
% very poor	0	33	0	N/A	0	0	N/A				
	Flov	Canala	Mustand	Couboss	Doc	l ontil	Chickers				
0/ overlent	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea				
% excellent	0	7	N/A	N/A	18	0	N/A				
% good	20	33	N/A	N/A	40	61	N/A				
% fair	80	55	N/A	N/A	37	39	N/A				
% poor	0	5	N/A	N/A	5	0	N/A				
% very poor	0	0	N/a	N/A	0	0	N/A				

Crop Districts and Rural Municipalities in Saskatchewan







Data Source:

Ν


Crop Districts - Saskatchewan Ministry of Agriculture

Geomatics Services, Ministry of Agriculture

April 9, 2018

Weekly Rainfall

from May 28 to June 3, 2019



Weekly Rainfall Summary

(in millimeters)

1 inch = 25 mm


for the period from May 28 to June 3, 2019

Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr	Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr	Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apı
1A	2	Mount Pleasant	NiL	56	4A	51	Reno	0.8	76.7	7A	287	St. Andrews	NiL	6
	3	Enniskillen	1	41		79 A	Arlington	NiL	53		288	Pleasant Valley	NiL	14
	32	Reciprocity	N/A	N/A		79 B	Arlington	NiL	92.5		290 A	Kindersley	N/A	9.4
	34	Browning	NiL	54		109	Carmichael	NiL	31.7		290 B	Kindersley	N/A	3
	61 64	Antler Brock	N/A NiL	26 51		110 111	Piapot Maple Creek	NiL N/A	52 NIL		292 317	Milton Marriott	NiL N/A	10
	65	Tecumseh	NiL	21	4B	139 A	Gull Lake	N/A	25		320 A	Oakdale	NiL	16.5
1B	91	Maryfield	NiL	30	70	139 B	Gull Lake	N/A	NIL		320 A	Oakdale	NiL	7
	94	Hazelwood	NiL	24.8		169	Pittville	N/A	27.5		321	Prairiedale	N/A	8
	122	Martin	NiL	41		229	Miry Creek	NiL	38.9	7B	347	Biggar	NiL	33
	123	Silverwood	NiL	34		231	Happyland	NiL	17		350	Mariposa	NiL	7
	124 A	Kingsley	1	31	5A	152	Spy Hill	18	45		351	Progress	NiL	17
		Kingsley	NiL	20.6		183	Fertile Belt	NiL	37		352	Heart's Hill	NiL	18
		Chester	N/A	39		211	Churchbridge	NiL	35		377	Glenside	NiL	15.5
		Chester	3	40		213	Saltcoats	TR	26		378	Rosemount	NiL	27
	151	Rocanville	7	45		217	Lipton	1	35		379	Reford	NiL	43
	154 A 154 B	Elcapo	TR	23 11		241 243	Calder Wallace	N/A	18		382	Eye Hill	NiL	27 27
	155	Elcapo Wolseley	6	47		243 244	Orkney	NiL NiL	19 18		409 A 409 B	Buffalo Buffalo	NiL NiL	23.4
2A	67	Weyburn	NiL	27		244 245 A	Garry	N/A	30		409 B	Round Valley	NiL	9.1
∠ /٦	68	Brokenshell	NiL	32		245 A 245 B	Garry	NiL	33	8A	394	Hudson Bay	10	55
	96	Fillmore	N/A	N/A		245 C	Garry	N/A	NIL	J, (395	Porcupine	NiL	27
	97	Wellington	2	35		246 A	Ituna Bon Accord	TR	29		397	Barrier Valley	NiL	28.8
2B	127 A	Francis	1.5	49		246 B	Ituna Bon Accord	9.7	40.1		426	Bjorkdale	8	33
	127 B	Francis	N/A	30.5		247	Kellross	NiL	23		427	Tisdale	NiL	5
	129	Bratt's Lake	N/A	33		248	Touchwood	NiL	18		428	Star City	6	28
	131	Baildon	NiL	37	5B	273	Sliding Hills	1	8		456	Arborfield	4	44.5
		Indian Head	0.3	39.5		277	Emerald	N/A	17		457 A	Connaught	NiL	16
		Indian Head	TR	64		305	Invermay	3	22		457 B	Connaught	N/A	15
	159	Sherwood	TR	37		307	Elfros	NiL	18		486	Moose Range	N/A	23
	160 A		N/A	28		308 A	Big Quill	NiL	10	op.	487	Nipawin	1	52.5
	160 B		TR	22		308 B	Big Quill	N/A	14	8B	369	St. Peter	TR	19
		Moose Jaw Moose Jaw	TR NiL	34 28		331 336	Livingston	N/A NiL	NIL 30		370 A 370 B	Humboldt Humboldt	TR 4	9 29
	162 A		NiL	24		337	Sasman Lakeview	N/A	7.5		370 B	Bayne	NiL	38
	162 B		N/A	53		366	Kelvington	TR	18.5		372	Grant	NiL	18.9
	191	Marquis	N/A	16		367	Ponass Lake	N/A	NIL		400	Three Lakes	NiL	38
ASE		Laurier	NiL	26.5	6A	190 A	Dufferin	N/A	27		429 A	Flett's Springs	TR	21
		Laurier	TR	36		190 B	Dufferin	N/A	16		429 B	Flett's Springs	TR	19
	39	The Gap	NiL	38		190 C	Dufferin	NiL	29		459	Kinistino	N/A	18
ASW	10	Happy Valley	NiL	35		190 D	Dufferin	N/A	17		460	Birch Hills	NiL	15.5
	12	Poplar Valley	N/A	39		219 A	Longlaketon	N/A	24	9AE	488	Torch River	16	58
	43	Old Post	NiL	58		219 B	Longlaketon	N/A	19.5		491	Buckland	5	49
		Stonehenge	NiL	41.9		220	McKillop	N/A	7		520	Paddockwood	N/A	26.5
		Stonehenge	NiL	40		221	Sarnia	TR	32.2		521	Lakeland	N/A	26.5
	74	Wood River	NiL	34.6		222	Craik	NiL	27	9AW	405	Great Bend	TR	3
BAN	102	Lake Johnston	N/A NiL	40 20		251	Big Arm	N/A NiL	9 19		406 A	Mayfield	2	15
	103	Sutton Hillsborough		-		252	Arm River Mount Hope		-		406 B 435	Mayfield Redberry	N/A 2	11
		Hillsborough	NiL NiL	31.5 50		279 282	McCraney	0.1 NiL	5.2 16		436	Douglas	NiL	26 28
	193	Eyebrow	N/A	21		313	Lost River	NiL	17		463	Duck Lake	N/A	31
BBS	17	Val Marie	NiL	31.8		339	Leroy	TR	28.8		466	Meeting Lake	TR	49
-	75	Pinto Creek	NiL	42		340	Wolverine	NiL	21		467 A	Round Hill	NiL	38
	77	Wise Creek	N/A	72		341	Viscount	NiL	16		467 B	Round Hill	2	39
	78	Grassy Creek	NiL	57.8		343	Blucher	NiL	21		493	Shellbrook	NiL	TR
	106	Whiska Creek	NII	17	6B	223 A	Huron	NiL	2		494	Canwood	N/A	5
	107	Lac Pelletier	N/A	31		223 B	Huron	NiL	22.5		497	Medstead	N/A	14.
	108	Bone Creek	NiL	30		284 A	Rudy	NiL	20.7	9B	438	Battle River	N/A	24
BN	138 A		NiL	27		284 B	Rudy	NiL	19		440	Hillsdale	NiL	39.
	138 B		N/A	NIL		285	Fertile Valley	NiL	25		442	Manitou Lake	NiL	18.
	165	Morse	N/A	11		286	Milden	NiL	35		498	Parkdale	NiL	28
		Riverside	NiL	34		314	Dundurn	NiL	32.5		499	Mervin	NiL	101.
		Riverside	N/A	26.5		344 A 344 B	Corman Park	NiL N/A	18		501 A		N/A	NII 40
	226	Victory	N/A Nii	2 15			Corman Park	N/A	10.6		501 B	Frenchman Butte	NiL N/A	40
	228 257	Lacadena Monet	NiL NiL	15 8		345 346	Vanscoy Perdue	NiL NiL	15 12		501 C 502	Frenchman Butte Britannia	N/A 2	26 6
	201	INIOLICE	INIL	U		346 376	Eagle Creek	NiL	22		561	Loon Lake	2	32
						403	Rosthern	NiL	28		588 A		3.5	29
						100		INIL	20		588 B		5	39
											622	Beaver River	N/A	25

These precipitation amounts represent point locations within each municipality and do not necessarily reflect the whole R. M.


Cumulative Rainfall

from April 1 to June 3, 2019




Cropland Topsoil Moisture Conditions

June 3, 2019




Hay and Pasture Topsoil Moisture Conditions June 3, 2019



Minimum Temperature

from May 28 to June 3, 2019



Maximum Temperature

from May 28 to June 3, 2019

