

West Nile Virus Surveillance Report, 2021: July 10

Table of Contents

1. West Nile virus transmission risk	page 2
2. Degree day accumulations	page 3
3. Mosquito surveillance results	page 5
4. West Nile virus animal cases	page 7
5. West Nile virus human cases	page 8

1. West Nile virus transmission risk (week ending July 10, 2021)

- *Culex tarsalis* mosquitoes continue to increase in southern Saskatchewan; to date, all mosquito pools have tested negative for WNV.
- High numbers of nuisance mosquito species continue to be detected in southeastern Saskatchewan.
- WNV transmission risk is currently minimal to low, but can increase later this season.
- To avoid being bitten, use mosquito repellents, cover up, and limit time outside during peak times of mosquito activity.

The risk of acquiring West Nile virus (WNV) infection in humans depends on various factors including time of year, number and location of infected *Culex tarsalis* mosquitoes, and number of days with sufficient heat. *Culex tarsalis* is a common summer mosquito throughout the agricultural portions of Saskatchewan, and is abundant in the southern areas of the province where it is hotter and drier. It is rarely found in the northern forested areas.

The risk of WNV transmission is low in the spring but often rises through the early and midsummer period, reaching a peak during the latter part of July and August. Infected, overwintered *Culex tarsalis* females may pose a small risk of transmission in spring.

The WNV risk levels may vary from minimal, when *Culex tarsalis* mosquitoes are rare and the weather has not been conducive for virus to cycle in mosquitoes and birds, to high when there are high numbers of WNV-infected mosquitoes and the weather and habitat conditions have been optimal for mosquito population growth, biting activity and transmission of the virus to humans.

Risk levels are determined in Saskatchewan through mosquito surveillance indicators such as *Culex tarsalis* numbers and infection rates, degree day or heat accumulation and other relevant weather factors such as precipitation. The level of risk in mosquitoes is determined by

using infection rates in mosquitoes (expressed as the number of infected mosquitoes/1000) and risk index calculated as: the infection rate X the average *Culex tarsalis* per trap night/1000).

Other relevant factors that help determine risk to humans include: time of year, the status of mosquito larval populations, past and predicted weather patterns, adult mosquito population age and trend, proximity to populated areas and other indicators such as positive birds or horses.

West Nile Virus Risk

Minimal - The mosquito species that carries WNV has not been detected. This does not mean risk is zero.

Low - The mosquito species that carries WNV has been detected in small numbers. There is a low probability of being bitten by an infected mosquito.


Moderate - WNV positive mosquitoes have been detected in numbers where there is a moderate probability of being bitten by an infected mosquito.

High - High numbers of WNV positive mosquitoes have been identified and are widespread. There is an increasing and high probability of being bitten by an infected mosquito.

2. Degree day accumulation

- The warmer temperatures of the past week have rapidly accelerated the rate of accumulation of degree days (Figure 1).
- In most parts of the province, 250-300 degree days have accumulated, which will support the emergence of the second generation of *Culex tarsalis*.

Figure 1: Degree day accumulations Prairie Provinces (July 11, 2021)



Note: The map covers the agricultural portion of Prairie Provinces. There is more habitat and higher numbers of *Culex tarsalis* in the southern portion of this area. *Culex tarsalis* is rarely found in forested areas. Map courtesy of Agriculture and Agri-food Canada.

Degree day: a measurement of heat accumulation from April 1. The threshold temperature below which WNV development and transmission is unlikely to occur in *Culex tarsalis* mosquitoes is 14.3°C. Degree days are calculated by subtracting the threshold or base temperature from the daily mean temperature. These are then summed to provide the total accumulation for the season.

Example: Mean daily temperature = 19.3°C; threshold temperature = 14.3°C; $19.3 - 14.3 = 5.0$ degree days.

Degree days are used in two ways. First, to predict *Culex tarsalis* development throughout the season by recording the total of accumulated degree days. On average, it takes approximately 250 to 300 degree days

(base 14.3° C) before the second generation of *Culex tarsalis* emerges. Females of this generation are most numerous and are largely responsible for transmission of WNV to humans. A total of 109 degree days are required for virus development to be completed within a particular population and for potential transmission to occur.

The second use of degree days is to determine the WNV transmission risk of infected mosquitoes. The risk of WNV transmission increases with increasing degree day accumulation. Moreover, consistently warmer temperatures will significantly shorten virus development time in the mosquitoes. This increases the potential risk of WNV transmission, if the virus is present and other conditions are favourable.

3. Mosquito surveillance results, 2021

- *Culex* spp. mosquitoes increased in southern Saskatchewan this week; however, numbers are much lower compared to the 5-year average (5 *Culex* spp. mosquitoes per trap night).
- To date, all mosquito pools have tested negative for WNV resulting in a low risk of transmission to humans.
- Compared to last year, higher numbers of nuisance mosquitoes were detected in southeastern Saskatchewan, likely due to increased precipitation recorded in the region since April.
- Warming nighttime temperatures will extend biting and egg-laying activities in the evening and early overnight periods.

Number of *Culex tarsalis* mosquitoes

Table 1: Average number of *Culex tarsalis* mosquitoes captured by date and ecological risk area, 2021

Surveillance Week ending	(1) Boreal Forest ^{‡§}	(2) Boreal Transition [§]	(3) Moist Mixed-Grass Prairie – Aspen Parkland	(4) Mixed-Grass Prairie
Jun 12	-	-	0	0
Jun 19	-	-	0	0
Jun 26	-	-	0.03	0
Jul 03	-	-	0.08	0
Jul 10	-	-	0.50	0.23
Jul 17	-	-		
Jul 24				
Jul 31				
Aug 07				
Aug 14				
Aug 21				
Aug 28				
Sep 04				
Sep 11				
Average*	-	-	0.12	0.05

Notes:

*Averages are determined by dividing the total number of *Culex tarsalis* mosquitoes caught by the total number of trapping nights.

‡ Due to extremely low numbers of *Culex* mosquitoes, trapping is generally not done in the Boreal Forest ecological risk area.

§ In 2021, trapping in the Boreal Forest and Transition ecological risk areas will start in mid-July if *Culex* spp. mosquitoes are detected in the southern traps by early July.

Number of mosquito pools testing positive

Table 2: Number of WNV positive mosquito pools *, percent positive pools and total number of pools tested by date and ecological risk area, 2021

Week Ending	(1) Boreal Forest ^{‡ §}			(2) Boreal Transition [§]			(3) Moist Mixed-grass Prairie-Aspen Parkland			(4) Mixed-grass Prairie			Weekly Totals		
	Positive	Tested	%	Positive	Tested	%	Positive	Tested	%	Positive	Tested	%	Positive	Tested	%
Jun 12	-	-	-	-	-	-	0	0	0	0	0	0	0	0	0
Jun 19	-	-	-	-	-	-	0	0	0	0	0	0	0	0	0
Jun 26	-	-	-	-	-	-	0	1	0	0	0	0	0	1	0
Jul 03	-	-	-	-	-	-	0	3	0	0	0	0	0	3	0
Jul 10	-	-	-	-	-	-	0	19	0	0	4	0	0	23	0
Jul 17	-	-	-	-	-	-									
Jul 24															
Jul 31															
Aug 07															
Aug 14															
Aug 21															
Aug 28															
Sep 04															
Sep 11															
Total	-	-	-	-	-	-	0	23	0	0	4	0	0	27	0

Notes:

* **Mosquito Pool** - Mosquitoes of the same species, collected from the same trap on the same date are pooled together for the purposes of laboratory testing. *Culex* mosquitoes (including *Culex tarsalis*, *Culex restuans* and *Culex territans*) collected from one trap on a given night are placed in pools of 1 - 50 mosquitoes for WNV testing. Other species, most notably *Culiseta inornata*, are occasionally placed in pools and tested as well. When more than 50 mosquitoes are collected from the same trap, multiple pools are tested. A positive pool refers to the detection of WNV in one or more mosquitoes collected from a given trap.

Percent positive pools are calculated as follows:

$$\frac{(\text{Number of positive pools})}{(\text{Total number tested})} \times 100 = \text{Percent positive pools}$$

[‡] Due to extremely low numbers of *Culex* mosquitoes, trapping is generally not done in the Boreal Forest ecological risk area.

[§] In 2021, trapping in the Boreal Forest and Boreal Transition ecological risk areas will start in mid-July if *Culex* spp. mosquitoes are detected in the southern traps by early July.

4. West Nile virus animal cases, 2021

Infections in animals such as horses are seasonal and often occur later in the season. The virus is well established in mosquito vectors in Saskatchewan. As WNV infections in horses lag behind infections in mosquitoes, mosquito surveillance provides more timely information about the risk to the public. Infections in horses can provide an indication that infections in humans may be occurring as well.

Table 3: Number of WNV positive horses by date and ecological risk area, July 04 2021 to the week ending July 10 2021

Week ending	(1) Boreal Forest	(2) Boreal Transition	(3) Moist Mixed-Grass Prairie-Aspen Parkland	(4) Mixed-Grass Prairie	Weekly totals
Jun 12	0	0	0	0	0
Jun 19	0	0	0	0	0
Jun 26	0	0	0	0	0
Jul 03	0	0	0	0	0
Jul 10	0	0	0	0	0
Jul 17					
Jul 24					
Jul 31					
Aug 07					
Aug 14					
Aug 21					
Aug 28					
Sep 04					
Sep 11					
Total	0	0	0	0	0

5. West Nile virus human cases, 2021 and 2003–2020

As with horses, human infections are seasonal and are often not detected until later in the season. Mosquito surveillance and other environmental risk indicators provide a more timely indication of risk.

Table 4: WNV surveillance in humans, July 04 2021 to the week ending July 10 2021

Number of WNV Positive Lab Tests*	WNV Neuroinvasive Disease ‡	WNV Deaths
0	0	0

Notes:

* These include tests done by the Roy Romanow Provincial Laboratory (RRPL) and Canadian Blood Services (CBS).

‡ The most useful indicator for the burden of disease in the general population is WNV neuroinvasive disease. For every case of WNV neuroinvasive disease, there are approximately 150 WNV infections in humans. The vast majority of people with WNV infections do not seek medical care.

Table 5: Saskatchewan Human WNV neuroinvasive cases 2003–2020*

Year	Neuroinvasive Cases	Deaths
2003	63	7
2004	0	0
2005	6	3
2006	3	0
2007	76	6
2008	1	0
2009	0	0
2010	0	0
2011	0	0
2012	0	0
2013	7	1
2014	1	0
2015	0	0
2016	0	0
2017	1	1
2018	3	1
2019	0	0
2020	0	0
Total	161	19

Note:

*Deaths are included in WNV Neuroinvasive disease case numbers except for 2003 when two deaths occurred in people with non-neuroinvasive West Nile Fever.