

Crop Report

For the Period August 24 to August 30, 2021

Published by the Ministry of Agriculture
ISSN 0701 7085
Report number 18, September 2, 2021

Despite the recent rainfall delays that many producers experienced, harvest continues to progress quickly this week. It has advanced substantially in the northern regions. Thirty-six per cent of the crop is now in the bin, up from 29 per cent last week and well over the five-year (2016-2020) average of 22 per cent. An additional 30 per cent of the crop is now swathed or ready to straight-cut.

The southwest region continues to have the most progress in the province with 47 per cent of the crop now combined. The southeast region has 42 per cent combined, the west-central 36 per cent, the east-central 25 per cent, the northeast 34 per cent and the northwest 21 per cent.

Ninety-nine per cent of the winter wheat, 83 per cent of the fall rye, 85 per cent of the lentils, 84 per cent of the field peas, 57 per cent of the mustard, 40 per cent of the durum, 11 per cent of the chickpeas, 36 per cent of the spring wheat and 11 per cent of the canola has now been combined. An additional 19 per cent of the canola and 16 per cent of the mustard is swathed or ready to straight-cut.

Many parts of the province experienced scattered rain showers this week, varying from trace amounts up to 77 mm in the Limerick area. The Macklin area received 51 mm, the Conquest area 40 mm, the Roblin area 35 mm, the Goodeve area 30 mm and the Rama area 28 mm.

Harvest Progress in SK Per cent Combined All Crops	
5 year avg. (2016-2020)	22
Aug 31/20	28
Sept 2/19	11
Aug 27/18	27
Aug 28/17	26
Aug 29/16	18
10-year avg. (2011-2020)	20

Recent rains have helped green up pastures which might allow cattle producers to continue pasture grazing and save their winter feed stocks.

One year ago

Twenty-eight per cent of the 2020 crop had been combined and 27 per cent was swathed or ready to straight-cut. Crops are drying down quickly and the majority of winter cereals and pulses have been combined. Lack of rain has allowed the topsoil moisture ratings to deteriorate.

Follow the 2021 Crop Report on Twitter @SKAgriculture.

Saskatchewan Harvest August 30, 2021 Per cent combined	
Winter wheat	99
Fall rye*	83
Spring wheat	36
Durum	40
Oats**	37
Barley***	47
Canaryseed	22
Flax	2
Canola	11
Mustard	57
Soybeans	34
Lentils	85
Peas	84
Chickpeas	11
*includes 10 per cent 'other'	
**includes three per cent 'other'	
***includes two per cent 'other'	

For further information, contact Matthew Struthers, AAg,
Crops Extension Specialist, Regional Services Branch,
Toll Free: 1-866-457-2377 or 306-694-3714, Email: cropreport@gov.sk.ca.
Also available on the Ministry of Agriculture website at saskatchewan.ca/crop-report.



Agriculture and
Agri-Food Canada

SCIC
SASKATCHEWAN CROP
INSURANCE CORPORATION

Saskatchewan

Topsoil moisture conditions continue to improve in most areas due to cooler temperatures and recent rains. Cropland topsoil moisture is rated as one per cent surplus, 40 per cent adequate, 34 per cent short and 25 per cent very short. Hay and pasture land topsoil moisture is rated as 28 per cent adequate, 43 per cent short and 29 per cent very short.

Most crop damage this past week was due to heavy rain, hail, strong winds, insects and the long-lasting effects of the drought. The rain has caused downgrading of many crops still standing in fields; the biggest issues reported are bleaching, staining, sprouting, low kernel weights and fungal growth. Reported yields are far lower than average for many parts of the province with some areas reporting some fields that have yielded almost nothing.

Producers are busy hauling bales and water as well as combining and swathing between rain showers.

Saskatchewan Harvest by Crop District August 30, 2021					
Crop District	Per cent combined	Crop District	Per cent combined	Crop District	Per cent combined
1A	48	4A	57	7A	32
1B	39	4B	72	7B	35
2A	45	5A	22	8A	32
2B	33	5B	20	8B	39
3ASE	66	6A	36	9AE	29
3ASW	35	6B	40	9AW	22
3AN	26			9B	20
3BS	56				
3BN	52				

As of Sept. 1, the AgriRecovery program, now referred to as the 2021 Canada-Saskatchewan Drought Response Initiative, is accepting producer applications. The Initiative will consist of two payments totalling up to \$200/head for cattle, with adjustments based on animal unit equivalents for other livestock. The initial payment will provide producers with \$100 per breeding female equivalent in inventory as of Aug. 1, 2021. Secondary payments of up to \$100 per breeding female in inventory as of Dec. 31, 2021, will be made to producers who have incurred additional costs to retain the animals. Producers with questions can call the initiative's dedicated toll-free number at 1-844-723-1211 or directly email skdri@scic.ca.

With harvest underway in Saskatchewan, we want to remind producers to exercise caution while working out in the field. Be aware, take breaks and remain safe.

The Farm Stress Line is also available for support 24-hours-a-day, seven-days-a-week, toll-free at 1-800-667-4442. Calls are answered by Mobile Crisis Services Regina, a non-profit, community-based agency and there is no call display.

Southeastern Saskatchewan:

- Crop District 1 – Carnduff, Estevan, Redvers, Moosomin and Kipling areas
- Crop District 2 – Weyburn, Milestone, Moose Jaw, Regina and Qu'Appelle areas
- Crop District 3ASE – Radville, Minton and Lake Alma areas

Even though there was scattered rainfall throughout the southeast region over the week, producers were able to get out into their fields and continue harvesting. Forty-two per cent of the crop is now in the bin, up from 30 per cent last week. The five-year (2016-2020) average for this time of year is 34 per cent. An additional 26 per cent of the crop is swathed or ready to straight cut. Rainy weather has resulted in downgrading of cereals and pulses but has given pastures a chance to recover from the season long drought.

Rainfall last week ranged from one mm to 74 mm in the Briercrest area. The Broadview and Glenavon areas reported 17 mm of rain, the Radville area 24 mm and the Weyburn area 32 mm.

Topsoil moisture conditions have slightly improved in the region with the recent rains. Cropland topsoil moisture is rated as 57 per cent adequate, 36 per cent short and seven per cent very short. Hay and pasture land topsoil moisture is rated as 27 per cent adequate, 60 per cent short and 13 per cent very short.

Crop damage this week was attributed to heavy rains, strong winds, insects and hail. One producer reported severe damage to as much as 1,500 acres of canola due to hail. Others have reported that hail blanketed the ground and gave the impression it had snowed. The total extent of the damage for the entire region is not known at this time but it is likely several fields were lost where the hail was the most severe.

Producers are starting to move cattle, hauling water and waiting for the conditions to dry up so they can resume harvest.

Southwestern Saskatchewan:

- Crop District 3ASW – Coronach, Assiniboia and Ogema areas
- Crop District 3AN – Gravelbourg, Mossbank, Mortlach and Central Butte areas
- Crop District 3B – Kyle, Swift Current, Shaunavon and Ponteix areas
- Crop District 4 – Consul, Maple Creek and Leader areas

Harvest progress was delayed greatly by rainfall this week, allowing only a small amount of crop to be harvested. Forty-seven per cent of the crop is now in the bin, up from 43 per cent last week but still well ahead of the five-year (2016-2020) average of 38 per cent. Still, many producers in the region have completed harvest.

The majority of the region did not receive significant amounts rain last week, although the Limerick area reported 77 mm. The Moose Jaw area reported 56 mm, the Mossebank area 27 mm, the Tyner area 13 mm and the Shaunavon area five mm.

Topsoil moisture conditions are slowly improving in the region thanks to several days of cool rainy weather. Cropland topsoil moisture is rated as 28 per cent adequate, 38 per cent short and 34 per cent very short. Hay and pasture land topsoil moisture is rated as 11

per cent adequate, 40 per cent short and 49 per cent very short. Crop District 4A reported that 63 per cent of the cropland and 67 per cent of the hay and pasture land is very short topsoil moisture at this time.

Most crop damage this past week was due to drought stress, strong winds, heavy rains and hail. Grasshoppers are still a large problem across the region and producers are trying to harvest as quickly as possible before more damage is caused, grasshoppers are also making grain cleaning a challenge.

Producers are busy moving bales and cattle while they wait for weather conditions to improve and let harvest resume.

East-Central Saskatchewan:

- Crop District 5 – Melville, Yorkton, Cupar, Kamsack, Foam Lake, Preeceville and Kelvington areas
- Crop District 6A – Lumsden, Craik, Watrous and Clavet areas

Very little harvest progress was made over the past week, only twenty-five per cent of the crop is now combined, up from 22 per cent last week and well ahead of the five-year (2016-2020) average of 13 per cent for this time of year. Now that it has rained and some of the crop is coming off tough, producers are using aerations bins or grain dryers to lower moisture levels.

Rainfall in the region ranged from trace amounts to 73 mm in the Rose Valley area. The Bethune area reported 39 mm of rain, the Roblin area 35 mm, the Rama area 28 mm and the Kenaston area 15 mm.

Topsoil moisture conditions have slightly improved in the region with the recent cool and rainy weather. Cropland topsoil moisture is rated as 60 per cent adequate, 24 per cent short and 16 per cent very short. Hay and pasture land topsoil moisture is rated as 53 per cent adequate, 31 per cent short and 16 per cent very short. Crop District 6A is reporting that 80 per cent of the cropland and 12 per cent of the hay and pasture land is adequate topsoil moisture at this time.

The majority of crop damage this week was due to strong winds, heavy rains and localized hail. The strong winds have blown swaths around and have lodged or shelled out some crops. The rainy humid conditions have resulted in bleaching, staining, sprouting and lowered kernel weights for many crops.

Producers are busy hauling bales, moving cattle and combining or swathing where it is drier.

West-Central Saskatchewan:

- Crop District 6B – Hanley, Outlook, Loreburn, Saskatoon and Arelee areas
- Crop District 7A – Rosetown, Kindersley, Eston and Major areas
- Crop District 7B – Kerrobert, Macklin, Wilkie and Biggar areas

The west-central region was able to get some harvesting done before being delayed by rains. Thirty-six per cent of the crop is now combined, up from 27 per cent last week and well ahead of the five-year (2016-2020) average of 17 per cent for this time of year. Many crops in the region that suffered severely from the drought were either cut for greenfeed or not cut at all. Crop quality issues such as lighter bushel weight and small, shriveled seed are very common in the region.

Much of the region received rain this past week with the Macklin area reporting 51 mm. The Kindersley area reported 47 mm, the Conquest area 40 mm, the Kerrobert area 12 mm and the Hanley area 11 mm.

Despite the incredible moisture deficit across the region, recent rain showers have been able to improve the topsoil moisture conditions in the region. Cropland topsoil moisture is rated as six per cent surplus, 28 per cent adequate, 42 per cent short and 24 per cent very short. Hay and pasture land topsoil moisture is rated as five per cent surplus, 21 per cent adequate, 37 per cent short and 37 per cent very short.

Drought conditions, strong winds and localized hail were the main causes of crop damage this past week. The strong winds have lodged crops, blown swaths around and shelled out standing crops. The rain has not slowed down the grasshoppers in the region and they still continue to cause issues for producers.

Producers are busy combining, swathings, hauling bales and getting ready to move cattle.

Northeastern Saskatchewan:

- Crop District 8 – Hudson Bay, Tisdale, Melfort, Carrot River, Humboldt, Kinistino, Cudworth and Aberdeen areas
- Crop District 9AE – Prince Albert, Choiceland and Paddockwood areas

Harvest is nicely progressing in the region despite recent rain and cooler weather. Thirty-four per cent of the crop is now in the bin, up from 25 per cent last week and ahead of the five-year (2016-2020) average of seven per cent for this time of year.

Overall crops remain in poor condition with yields expected to be below average. While some crops are coming off dry, others are coming off tough and are being placed into aeration bins and dryers. Some fields remain wet and will need good drying weather before harvest can begin.

Most of the region received rain last week that delayed some producers in the field. The Melfort area reported 56 mm, the Humboldt area 53 mm, the Prince Albert area 50 mm, the Lake Lenore area 30 mm and the Hudson Bay and Nipawin areas 22 mm.

Cropland topsoil moisture is rated as 42 per cent adequate, 47 per cent short and 11 per cent very short. Hay and pasture land topsoil moisture is rated as 26 per cent adequate, 58 per cent short and 16 per cent very short. Crop District 8A is reporting that 58 per cent of the cropland and 57 per cent of the hay and pasture land is short for topsoil moisture at this time.

The majority of crop damage this past week was due to localized hail and strong winds that have blown swaths around and shelled some standing crops. One hail storm was reported to have covered approximately 20 quarters and the damage varied from light to 100 per cent loss.

Producers are busy combining, swathing, desiccating and hauling bales.

Northwestern Saskatchewan:

- Crop District 9AW – Shellbrook, North Battleford, Big River and Hafford areas
- Crop District 9B – Meadow Lake, Turtleford, Pierceland, Maidstone and Lloydminster areas

The majority of the northwest did not receive large amounts of rainfall but conditions were still damp enough to slow down harvest progress across the region. Twenty-one per cent of the crop is now combined, up from 15 per cent last week and more than four times the five-year (2016-2020) average of five per cent. Initial crop yields and qualities are very low.

Rainfall this past week ranged from trace amounts to 50 mm in the Duck Lake area. The Hafford and Barthel areas reported 10 mm of rain while the Mayfair and North Battleford areas reported four mm.

Cropland topsoil moisture conditions are rated as 13 per cent adequate, 34 per cent short and 53 per cent very short. Hay and pasture land topsoil moisture is rated as eight per cent adequate, 36 per cent short and 56 per cent very short.

Most crop damage this past week was due to strong winds that have blown swaths around and shelled out some standing crops. Hail was reported in some areas but the damage is not known at this time.

Producers are busy combining, swathing, desiccating and hauling bales.


Saskatchewan Harvest Progress - August 30, 2021

*Other - crop that will not be harvested due to weather, insect or disease damage or will be greenfeed or silage

Winter Wheat	% Standing	% in swath	% ready to straight combine	% combined	% other (greenfeed/silage)
southeast	0	0	0	99	1
southwest	0	0	0	99	1
east central	0	0	0	100	0
west central	0	0	0	98	2
northeast	0	0	0	100	0
northwest	n/a	n/a	n/a	n/a	n/a
provincial	0	0	0	99	1
Fall Rye	% Standing	% in swath	% ready to straight combine	% combined	% other (greenfeed/silage)
southeast	0	0	0	79	21
southwest	0	0	0	75	25
east central	0	0	0	90	10
west central	0	0	0	88	12
northeast	0	0	0	94	6
northwest	0	0	0	70	30
provincial	0	0	0	83	17
Spring Wheat	% Standing	% in swath	% ready to straight combine	% combined	
southeast	26	2	24	48	
southwest	24	5	31	40	
east central	36	8	25	31	
west central	30	8	30	32	
northeast	17	3	39	41	
northwest	40	5	25	30	
provincial	25	5	34	36	
Durum	% Standing	% in swath	% ready to straight combine	% combined	
southeast	22	5	27	46	
southwest	22	10	29	39	
east central	38	6	26	30	
west central	44	3	18	35	
northeast	N/A	N/A	N/A	N/A	
northwest	N/A	N/A	N/A	N/A	
provincial	25	8	27	40	
Barley	% Standing	% in swath	% ready to straight combine	% combined	% other (greenfeed/silage)
southeast	17	9	11	50	13
southwest	18	7	15	41	19
east central	28	7	15	47	3
west central	18	4	22	38	18
northeast	10	4	18	68	0
northwest	40	5	11	31	13
provincial	25	6	16	47	6
Oats	% Standing	% in swath	% ready to straight combine	% combined	% other (greenfeed/silage)
southeast	25	0	10	48	17
southwest	25	3	14	23	35
east central	46	0	20	24	10
west central	19	0	8	22	51
northeast	21	1	34	44	0
northwest	49	4	8	20	19
provincial	30	5	18	37	10
Canaryseed	% Standing	% in swath	% ready to straight combine	% combined	
southeast	64	0	31	5	
southwest	56	0	11	33	
east central	70	0	26	4	
west central	30	0	34	36	
northeast	85	0	11	4	
northwest	N/A	N/A	N/A	N/A	
provincial	70	0	8	22	


Flax	% Standing	% in swath	% ready to straight combine	% combined	
southeast	93	5	1	1	
southwest	91	0	3	6	
east central	84	3	12	1	
west central	80	0	18	2	
northeast	86	1	3	10	
northwest	84	0	12	4	
provincial	88	3	7	2	
Canola	% Standing	% in swath	% ready to straight combine	% combined	
southeast	39	17	25	19	
southwest	50	0	26	24	
east central	44	37	14	5	
west central	40	24	25	11	
northeast	40	32	19	9	
northwest	33	51	12	4	
provincial	41	29	19	11	
Mustard	% Standing	% in swath	% ready to straight combine	% combined	
southeast	15	5	36	44	
southwest	16	1	16	67	
east central	61	9	30	0	
west central	72	0	11	17	
northeast	N/A	N/A	N/A	N/A	
northwest	N/A	N/A	N/A	N/A	
provincial	24	3	16	57	
Soybeans	% Standing	% in swath	% ready to straight combine	% combined	
southeast	55	0	5	40	
southwest	40	0	10	50	
east central	63	0	7	30	
west central	28	0	55	17	
northeast	N/A	N/A	N/A	N/A	
northwest	N/A	N/A	N/A	N/A	
provincial	64	0	2	34	
Field Peas	% Standing	% in swath	% ready to straight combine	% combined	
southeast	0	0	1	99	
southwest	9	0	4	87	
east central	12	4	4	80	
west central	6	0	13	81	
northeast	3	0	3	94	
northwest	4	4	8	84	
provincial	11	1	4	84	
Lentils	% Standing	% in swath	% ready to straight combine	% combined	
southeast	1	0	5	94	
southwest	12	0	6	82	
east central	1	2	3	94	
west central	7	0	7	86	
northeast	17	1	20	62	
northwest	18	0	27	55	
provincial	8	1	6	85	
Chickpeas	% Standing	% in swath	% ready to straight combine	% combined	
southeast	56	0	6	38	
southwest	69	0	11	20	
east central	30	0	20	50	
west central	N/A	N/A	N/A	N/A	
northeast	N/A	N/A	N/A	N/A	
northwest	N/A	N/A	N/A	N/A	
provincial	64	0	25	11	

Crop Districts and Rural Municipalities in Saskatchewan



Weekly Rainfall

from August 24 to August 30, 2021



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Weekly Rainfall Summary

(in millimeters)

1 inch = 25 mm

for the period from August 24 to 30, 2021


Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr	Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr	Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr
1A	2	Mount Pleasant	4	201	4A	51	Reno	2.3	154.4	7A	287	St. Andrews	N/A	68
	3	Enniskillen	1	268		79 A	Arlington	3	153		288	Pleasant Valley	3	108
	32	Reciprocity	N/A	172		79 B	Arlington	TR	135		290	Kindersley	N/A	119.1
	61	Antler	N/A	340		109	Carmichael	N/A	78.6		292	Milton	N/A	174
	64	Brock	6	273		110	Piapot	N/A	69		317	Marriott	5	141
	65	Tecumseh	N/A	234		139	Gull Lake	N/A	81		320 A	Oakdale	47	177
1B	95	Golden West	N/A	56	4B	229	Miry Creek	N/A	96		320 B	Oakdale	19	126
	122	Martin	25	289		231	Happyland	N/A	105		321	Prairiedale	7	168
	123	Silverwood	24	276		181	Langenburg	N/A	132	7B	347	Biggar	5	190
	124	Kingsley	17	223	5A	183 A	Fertile Belt	33	295		350	Mariposa	12	79
	125 A	Chester	20	295		183 B	Fertile Belt	26	115		351	Progress	4	194
	125 B	Chester	17	273		211	Churchbridge	27	270		377	Glenside	23.5	146.5
	151	Rocanville	10	282		213	Saltcoats	27	268		378	Rosemount	N/A	236
	154 A	Elcapo	5	249		217	Lipton	24	305		382	Eye Hill	51	236.5
	154 B	Elcapo	N/A	38		241	Calder	35	278		409 A	Buffalo	N/A	111
	155	Wolseley	N/A	314		243	Wallace	NIL	257		409 B	Buffalo	22	173
2A	67	Weyburn	8	230.2		245 A	Garry	17	196		410	Round Valley	2	102.3
	68	Brokenshell	32	340.5		245 B	Garry	30	238	8A	394	Hudson Bay	22	176
	97	Wellington	3	308		246 A	Ituna Bon Accord	4	246		395	Porcupine	N/A	194
	100	Elmsthorpe	23	275.8		246 B	Ituna Bon Accord	N/A	186.6		397	Barrier Valley	24	156.4
2B	127 A	Francis	2.5	366.5		247	Kellross	10	237.1		428	Star City	20	116
	127 B	Francis	N/A	194		248	Touchwood	8	187		456	Arborkfield	13	176
	129	Bratt's Lake	6	320	5B	273	Sliding Hills	N/A	259.5		457	Connaught	N/A	92.7
	131	Baldon	74	306		277	Emerald	25	255		486	Moose Range	22	162
	156 A	Indian Head	16.3	275.2		301	St.Philips	N/A	160.6		487	Nipawin	N/A	88
	156 B	Indian Head	22	388		305	Invermay	28	238	8B	369	St. Peter	53	186
	159	Sherwood	TR	239		307	Elfros	N/A	197		370 A	Humboldt	19	154
	160	Pense	N/A	219		308 A	Big Quill	5	166		370 B	Humboldt	N/A	61
	161 A	Moose Jaw	37	319		308 B	Big Quill	N/A	95		371	Bayne	42	175
	161 B	Moose Jaw	N/A	408		331	Livingston	16.5	236.3		372	Grant	27.6	152.7
	162 A	Caron	22.9	273.2		334	Preeceville	N/A	87		400	Three Lakes	30	143
	162 B	Caron	39	259		336	Sasman	17	198		429	Flett's Springs	46	143
	191	Marquis	N/A	144		337	Lakeview	N/A	138		430	Invergordon	56	187
3ASE	38 A	Laurier	24	179.4		366	Kelvington	28.5	248.5		459	Kinistino	50	190
	38 B	Laurier	24	281		367	Ponass Lake	73	231		460	Birch Hills	44.3	104.8
3ASW	10	Happy Valley	3	379	6A	190 A	Dufferin	39	271	9AE	461	Prince Albert	50	166
	43	Old Post	3	198		190 B	Dufferin	N/A	181		488	Torch River	35	132
	73 A	Stonehenge	28	186		190 C	Dufferin	38	332		491	Buckland	20	152
	73 B	Stonehenge	77	219.5		190 D	Dufferin	N/A	84		520	Paddockwood	N/A	241
	74	Wood River	28	245.5		219 A	Longlaketon	12	226		521	Lakeland	N/A	216
3AN	102	Lake Johnston	27	313.7		219 B	Longlaketon	N/A	247.2	9AW	405	Great Bend	N/A	44
	103	Sutton	N/A	117		220	McKillop	N/A	255		435	Redberry	10	161
	132 A	Hillsborough	22	254.5		221	Sarnia	24.2	124.6		436	Douglas	NIL	106
	132 B	Hillsborough	56	468		222	Craik	25	205		437	North Battleford	46	158.4
	193	Eyebrow	17	177		251	Big Arm	32.5	128		463	Duck Lake	50	192
3BS	75	Pinto Creek	4.5	160.5		252	Arm River	26	195		466	Meeting Lake	4	164
	77	Wise Creek	7	151		279	Mount Hope	N/A	209		467 A	Round Hill	4	152
	78	Grassy Creek	5	150.5		282	McCraney	15	136.8		467 B	Round Hill	4	164
	106	Whiska Creek	N/A	146		339	Leroy	12.2	190.8		493	Shellbrook	N/A	100
	107	Lac Pelletier	N/A	100		340	Wolverine	N/A	114.3		496	Spiritwood	N/A	175
	108	Bone Creek	N/A	151		341	Viscount	25	123		497	Medstead	5	76
3BN	138	Webb	7	141.5		343	Blucher	4	159	9B	440	Hillsdale	0.5	133
	165	Morse	4	165.8	6B	223 A	Huron	17	148		442	Manitou Lake	2	153.9
	168 A	Riverside	NIL	109		223 B	Huron	22	192.5		498	Parkdale	NIL	135.5
	168 B	Riverside	N/A	93.5		223 C	Huron	N/A	91.4		499	Mervin	NIL	175.9
	226	Victory	N/A	73		284 A	Rudy	NIL	107		501 A	Frenchman Butte	1	174.2
	228	Lacadena	13	108		284 B	Rudy	NIL	64		501 B	Frenchman Butte	NIL	164
	257	Monet	N/A	84.5		285	Fertile Valley	40	142.5		501 C	Frenchman Butte	2	124
						286	Milden	4	99		502	Britannia	N/A	58.5
						314	Dundurn	11	138		561	Loon Lake	10	158
						344	Corman Park	8	62		588	Meadow Lake	NIL	164
						376	Eagle Creek	11	133		622	Beaver River	NIL	117.5
						403	Rosthern	9	155					

Municipality No: A, B, C and D - more than one reporter

These precipitation amounts represent point locations within each municipality and do not necessarily reflect the whole R. M.


Cumulative Rainfall

from April 1 to August 30, 2021



Cropland Topsoil Moisture Conditions


August 30, 2021



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Hay and Pasture Topsoil Moisture Conditions


August 30, 2021



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Minimum Temperature


from August 24 to August 30, 2021



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Maximum Temperature

from August 24 to August 30, 2021



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.